

# 1 UNIT

## Set Theory, Functions and Natural Numbers

### CONTENTS

<b>Part-1</b>	: Set Theory : Introduction, ..... Combination of Sets, Multisets	1-2C to 1-5C
<b>Part-2</b>	: Ordered Pair, Proof of Some ..... General Identities on Sets	1-5C to 1-8C
<b>Part-3</b>	: Relations : Definition, Operation ..... on Relations, Properties of Relation, Composite Relation, Equality of Relation, Recursive Definition of Relation, Order of Relation	1-8C to 1-16C
<b>Part-4</b>	: Function : Definition, ..... Classification of Functions, Operations on Functions, Recursively Defined Functions, Growth of Functions	1-16C to 1-22C
<b>Part-5</b>	: Natural Numbers : ..... Introductions, Mathematical Induction, Variants of Induction, Induction with Non-zero Base Cases	1-22C to 1-26C
<b>Part-6</b>	: Proof Methods, Proof by ..... Counter-Example, Proof by Contradiction	1-26C to 1-29C

1-1 C (CS/IT-Sem-3)

## PART-1

*Set Theory : Introduction, Combination of Sets, Multisets.*

## Questions-Answers

## Long Answer Type and Medium Answer Type Questions

**Que 1.1.** What do you understand by set? Explain different types of set.

**Answer**

1. A set is a collection of well defined objects, called elements or members of the set.
2. These elements may be anything like numbers, letters of alphabets, points etc.
3. Sets are denoted by capital letters and their elements by lower case letters.
4. If an object  $x$  is an element of set  $A$ , we write it as  $x \in A$  and read it as ' $x$  belongs to  $A$ ' otherwise  $x \notin A$  ( $x$  does not belong to  $A$ ).

**Types of set :**

1. **Finite set :** A set with finite or countable number of elements is called finite set.
2. **Infinite set :** A set with infinite number of elements is called infinite set.
3. **Null set :** A set which contains no element at all is called a null set. It is denoted by  $\phi$  or  $\{\}$ . It is also called empty or void set.
4. **Singleton set :** A set which has only one element is called singleton set.
5. **Subset :** Let  $A$  and  $B$  be two sets, if every elements of  $A$  also belongs to  $B$  i.e., if every element of set  $A$  is also an element of set  $B$ , then  $A$  is called subset of  $B$  and it is denoted by  $A \subseteq B$ .  
Symbolically,  $A \subseteq B$  if  $x \in A \Rightarrow x \in B$ .
6. **Superset :** If  $A$  is subset of a set  $B$ , then  $B$  is called superset of  $A$ .
7. **Proper subset :** Any subset  $A$  is said to be proper subset of another set  $B$ , if there is at least one element of  $B$  which does not belong to  $A$ , i.e., if  $A \subseteq B$  but  $A \neq B$ . It is denoted by  $A \subset B$ .
8. **Universal set :** In many applications of sets, all the sets under consideration are considered as subsets of one particular set. This set is called universal set and is denoted by  $U$ .
9. **Equal set :** Two set  $A$  and  $B$  are said to be equal if every element of  $A$  belong to set  $B$  and every element of  $B$  belong to set  $A$ . It is written as  $A = B$ .  
Symbolically,  $A = B$  if  $x \in A$  and  $x \in B$ .

10. **Disjoint set** : Let  $A$  and  $B$  be two sets, if there is no common element between  $A$  and  $B$ , then they are said to be disjoint.

**Que 1.2.** Describe the different types of operation on sets.

**Answer**

- 1. Union** : Let  $A$  and  $B$  be two sets, then the union of sets  $A$  and  $B$  is a set that contain those elements that are either in  $A$  or  $B$  or in both. It is denoted by  $A \cup B$  and is read as 'A union B'.

Symbolically,  $A \cup B = \{x | x \in A \text{ or } x \in B\}$

**For example** :  $A = \{1, 2, 3, 4\}$   
 $B = \{3, 4, 5, 6\}$   
 $A \cup B = \{1, 2, 3, 4, 5, 6\}$
- 2. Intersection** : Let  $A$  and  $B$  be two sets, then intersection of  $A$  and  $B$  is a set that contain those elements which are common to both  $A$  and  $B$ . It is denoted by  $A \cap B$  and is read as 'A intersection B'.

Symbolically,  $A \cap B = \{x | x \in A \text{ and } x \in B\}$

**For example** :  $A = \{1, 2, 3, 4\}$   
 $B = \{2, 4, 6, 7\}$   
 then  $A \cap B = \{2, 4\}$
- 3. Complement** : Let  $U$  be the universal set and  $A$  be any subset of  $U$ , then complement of  $A$  is a set containing elements of  $U$  which do not belong to  $A$ . It is denoted by  $A^c$  or  $A'$  or  $\bar{A}$ .

Symbolically,  $A^c = \{x | x \in U \text{ and } x \notin A\}$

**For example** :  $U = \{1, 2, 3, 4, 5, 6\}$   
 and  $A = \{2, 3, 5\}$   
 then  $A^c = \{1, 4, 6\}$
- 4. Difference of sets** : Let  $A$  and  $B$  be two sets. Then difference of  $A$  and  $B$  is a set of all those elements which belong to  $A$  but do not belong to  $B$  and is denoted by  $A - B$ .

Symbolically,  $A - B = \{x | x \in A \text{ and } x \notin B\}$

**For example** : Let  $A = \{2, 3, 4, 5, 6, 7\}$   
 and  $B = \{4, 5, 7\}$   
 then  $A - B = \{2, 3, 6\}$
- 5. Symmetric difference of set** : Let  $A$  and  $B$  be two sets. The symmetric difference of  $A$  and  $B$  is a set containing all the elements that belong to  $A$  or  $B$  but not both. It is denoted by  $A \oplus B$  or  $A \Delta B$ .

Also  $A \oplus B = (A \cup B) - (A \cap B)$

**For example** : Let  $A = \{2, 3, 4, 6\}$   
 $B = \{1, 2, 5, 6\}$   
 then  $A \oplus B = \{1, 3, 4, 5\}$

**Que 1.3.** What do you mean by multisets ?

**Answer**

1. Multisets are sets where an element appear more than once

For example :  $A = \{1, 1, 1, 2, 2, 3\}$   
 $B = \{a, a, a, b, b, b, c, c\}$

are multisets.

2. The multisets  $A$  and  $B$  can also be written as  
 $A = \{3.1, 2.2, 1.3\}$  and  $B = \{3.a, 3.b, 2.c\}$
3. The multiplicity of an element in a multiset is defined to be the number of times an element appears in the multiset. In above examples, multiplicities of the elements 1, 2, 3 in multiset  $A$  are 3, 2, 1 respectively.
4. Let  $A$  and  $B$  be two multisets. Then,  $A \cup B$ , is the multiset where the multiplicity of an element is the maximum of its multiplicities in  $A$  and  $B$ .
5. The intersection of  $A$  and  $B$ ,  $A \cap B$ , is the multiset where the multiplicity of an element is the minimum of its multiplicities in  $A$  and  $B$ .
6. The difference of  $A$  and  $B$ ,  $A - B$ , is the multiset where the multiplicity of an element is equal to the multiplicity of the element in  $A$  minus the multiplicity of the element in  $B$  if the difference is positive, and is equal to zero if the difference is zero and negative.
7. The sum of  $A$  and  $B$ ,  $A + B$ , is the multiset where the multiplicity of an element is the sum of multiplicities of the elements in  $A$  and  $B$ .

**Que 1.4.** Let  $P$  and  $Q$  be two multisets  $\{4.a, 3.b, 1.c\}$  and  $\{3.a, 3.b,$

$2.d\}$  respectively. Find

- i.  $P \cup Q$ , ii.  $P \cap Q$ , iii.  $P - Q$ , iv.  $Q - P$ , v.  $P + Q$ .

**Answer**

- i.  $P \cup Q = \{4.a, 3.b, 1.c, 2.d\}$
- ii.  $P \cap Q = \{3.a, 3.b\}$
- iii.  $P - Q = \{1.a, 1.c\}$
- iv.  $Q - P = \{2.d\}$
- v.  $P + Q = \{7.a, 6.b, 1.c, 2.d\}$

**Que 1.5.** Describe each of following in roster form :

- i.  $A = \{x : x \text{ is an even prime}\}$
- ii.  $B = \{x : x \text{ is a positive integral divisor of } 60\}$
- iii.  $C = \{x \in R : x^2 - 1 = 0\}$
- iv.  $D = \{x : x^2 - 2x + 1 = 0\}$
- v.  $E = \{x : x \text{ is multiple of } 3 \text{ or } 5\}$

**Answer**

- i.  $A = \{2\}$
- ii.  $B = \{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$
- iii.  $C = \{1, -1\}$
- iv.  $D = \{1\}$
- v.  $E = \{3, 5, 6, 10, 15, \dots\}$

**Que 1.6.** Describe the following in set builder form :

- i.  $A = \{-4, -3, -2, -1, 0, 1, 2, 3\}$

- ii.  $B = \{1, 8, 27, 64\}$   
 iii.  $C = \{3, 6, 9, 12, 15, \dots\}$   
 iv.  $D = \{2, 3, 5, 7, 11, 13, \dots\}$

**Answer**

- i.  $A = \{x \mid x \text{ is an integer and } -4 \leq x \leq 3\}$   
 ii.  $B = \{x \mid x = n^3, \text{ where } 1 \leq n \leq 4, n \text{ is natural number}\}$   
 iii.  $C = \{x \mid x = 3n, \text{ where } n \text{ is natural number}\}$   
 iv.  $D = \{x \mid x \text{ is the integer and prime number}\}$

**PART-2***Ordered Pair, Proof of Some General Identities on Sets.***Questions-Answers****Long Answer Type and Medium Answer Type Questions**

**Que 1.7.** List down laws of algebra of sets.

OR

Write down the general identities on sets.

**Answer**

Let  $A, B, C$  be any three sets and  $U$  be the universal set, then following are the laws of algebra of sets :

1. **Idempotent laws :**
  - a.  $A \cup A = A$
  - b.  $A \cap A = A$
2. **Commutative laws :**
  - a.  $A \cup B = B \cup A$
  - b.  $A \cap B = B \cap A$
3. **Associative laws :**
  - a.  $A \cup (B \cap C) = (A \cup B) \cap C$
  - b.  $A \cap (B \cup C) = (A \cap B) \cup C$
4. **Distributive laws :**
  - a.  $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
  - b.  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
5. **Identity laws :**
  - a.  $A \cup \phi = A$
  - b.  $A \cap U = A$
  - c.  $A \cup U = U$
  - d.  $A \cap \phi = \phi$
6. **Involution law :**
  - a.  $(A^c)^c = A$

**7. Complement laws :**

- a.  $A \cup A^c = U$
- b.  $A \cap A^c = \phi$
- c.  $U^c = \phi$
- d.  $\phi^c = U$

**8. De Morgan's laws :**

- a.  $(A \cup B)^c = A^c \cap B^c$
- b.  $(A \cap B)^c = A^c \cup B^c$

**9. Absorption laws :**

- a.  $A \cup (A \cap B) = A$
- b.  $A \cap (A \cup B) = A$

**Que 1.8.** Prove for any two sets  $A$  and  $B$  that,  $(A \cup B)^c = A^c \cap B^c$ .

AKTU 2014-15, Marks 05

**Answer**

Let  $x \in (A \cup B)^c$   
 $\Rightarrow x \notin A \cup B$   
 $\Rightarrow x \notin A$  and  $x \notin B$   
 $\Rightarrow x \in A^c$  and  $x \in B^c$   
 $\Rightarrow x \in A^c \cap B^c$   
 $\Rightarrow (A \cup B)^c \subseteq A^c \cap B^c$  ... (1.8.1)

Now, let  $x \in A^c \cap B^c$   
 $\Rightarrow x \in A^c$  and  $x \in B^c$   
 $\Rightarrow x \notin A$  and  $x \notin B$   
 $\Rightarrow x \in (A \cup B)^c$   
 $\Rightarrow x \in (A \cup B)^c$   
 $\Rightarrow (A^c \cap B^c) \subseteq (A \cup B)^c$  ... (1.8.2)

From eq. (1.8.1) and (1.8.2),  $(A \cup B)^c = A^c \cap B^c$

**Que 1.9.** Let  $U = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$ , and the ordering of elements of  $U$  has the elements in increasing order; that is,  $a_i = i$ . What bit strings represent the subset of all odd integers in  $U$ , the subset of all even integers in  $U$ , and the subset of integers not exceeding 5 in  $U$ ?

**Answer**

Since  $U = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$

Let the subset of all odd integers be  $s_1$ , i.e.,

$$s_1 = \{1, 3, 5, 7, 9\}$$

subset of all even integers be  $s_2$ , i.e.,

$$s_2 = \{2, 4, 6, 8, 10\}$$

and the subset of integers not exceeding 5 be  $s_3$ , i.e.,

$$s_3 = \{1, 2, 3, 4, 5\}$$

The bit string by characteristic function is given as follows :

Bit string for  $s_1$  is 1010101010

Bit string for  $s_2$  is 0101010101

Bit string for  $s_3$  is 1111100000

**Que 1.10.** If  $A$  and  $B$  are two subsets of universal set, then prove the following :

a.  $(A - B) = (B - A) \text{ iff } A = B$

b.  $(A - B) = A \text{ iff } A \cap B = \phi$

**Answer**

a. Let  $A = B$   
 Consider any element  $x \in A - B$   
 $\Rightarrow x \in A$  and  $x \notin B$   
 $\Rightarrow x \in B$  and  $x \notin A$  [ $\because A = B$ ]  
 $\Rightarrow x \in B - A$   
 $\therefore A - B \subseteq B - A$  ... (1.10.1)  
 Conversely, if  $x \in B - A$   
 $\Rightarrow x \in B$  and  $x \notin A$   
 $\Rightarrow x \in A$  and  $x \notin B$   
 $\Rightarrow x \in A - B$   
 $B - A \subseteq A - B$  ... (1.10.2)

From eq. (1.10.1) and (1.10.2), we have

If  $A = B \Rightarrow A - B = B - A$

Now let  $A - B = B - A$

Let  $x \in A - B$  as  $A - B = B - A$

$\therefore x \in B - A$

Now  $x \in A - B \Rightarrow x \in A$  and  $x \notin B$  ... (1.10.3)

and  $x \in B - A \Rightarrow x \in B$  and  $x \notin A$  ... (1.10.4)

Eq. (1.10.3) and (1.10.4), can hold true when  $A = B$

b. Let  $A - B = A$   
 To show  $A \cap B = \phi$   
 Let  $A \cap B \neq \phi$  and let  $x \in A \cap B$  and  $x \notin \phi$   
 $\Rightarrow x \in A$  and  $x \in B$   
 $\Rightarrow x \in (A - B)$  and  $x \notin B$  [ $\because A - B = A$ ]  
 $\Rightarrow x \in A$  and  $x \notin B$  and  $x \in B$   
 $\Rightarrow x \in \phi$

which is a contradiction.

$\therefore A \cap B = \phi$

Now conversely, let  $A \cap B = \phi$

To show  $A - B = A$

Let  $x \in A - B$   
 $\Rightarrow x \in A$  and  $x \notin B$   
 $\Rightarrow x \in A$  [as  $A \cap B = \phi$ ]  
 $\Rightarrow A - B \subseteq A$  ... (1.10.5)

Conversely, let  $x \in A$

$\Rightarrow x \in A$  and  $x \notin B$  [as  $A \cap B = \phi$ ]

$$\begin{aligned} \Rightarrow & x \in A - B \\ \therefore & A \subseteq A - B \\ \text{From eq. (1.10.5) and (1.10.6),} & \\ & A = A - B \end{aligned} \quad \dots(1.10.6)$$

### PART-3

*Relations : Definition, Operation on Relations, Properties of Relation, Composite Relation, Equality of Relation, Recursive Definition of Relation, Order of Relation.*

### Questions-Answers

#### Long Answer Type and Medium Answer Type Questions

**Que 1.11.** Describe the term relation along with its types.

Let  $A$  and  $B$  be two non-empty sets, then  $R$  is relation from  $A$  to  $B$  if  $R$  is subset of  $A \times B$  and is set of ordered pair  $(a, b)$  where  $a \in A$  and  $b \in B$ . It is denoted by  $aRb$  and read as "a is related to b by R".

Symbolically,  $R = \{(a, b) : a \in A, b \in B, aRb\}$

If  $(a, b) \in R$  then  $a \not R b$  and read as "a is not related to b by R".

**For example :**

Let  $A = \{1, 2, 3, 4\}$ ,  $B = \{1, 2\}$  and  $aRb$  iff  $a \times b = \text{even number}$

Then  $R = \{(1, 2), (2, 1), (2, 2), (3, 2), (4, 1), (4, 2)\}$

**Types of relation :**

- 1. Universal relation :** A relation  $R$  is called universal relation on  $A$  if  $R = A \times A$ . In case where  $R$  is defined from  $A$  to  $B$ , then  $R$  is universal relation if  $R = A \times B$ .

**For example :**

If  $A = \{1, 2, 3\}$

then  $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3), (3, 1), (3, 2), (3, 3)\}$

is universal relation over  $A$ .

- 2. Identity relation :** A relation  $R$  is called identity relation on  $A$  if  $R = \{(a, a) | a \in A\}$ . It is denoted by  $I_A$  or  $\Delta_A$  or  $\Delta$ . It is also called diagonal relation.

**For example :**

If  $A = \{1, 2, 3\}$ , then  $I_A = \{(1, 1), (2, 2), (3, 3)\}$

is identity relation on  $A$ .

- 3. Void relation :** A relation  $R$  is called a void relation on  $A$  if  $R = \phi$ . It is also called null relation.

**For example :**

If  $A = \{1, 2, 3\}$  and  $R$  is defined as  $R = \{(a, b) | a + b > 5\}$ ,  $a, b \in A$  then  $R = \phi$ .

4. **Inverse relation** : A relation  $R$  defined from  $B$  to  $A$  is called inverse relation of  $R$  defined from  $A$  to  $B$  if

$$R^{-1} = \{(b, a) : b \in B \text{ and } a \in A \text{ and } (a, b) \in R\}.$$

**For example** : Consider relation

$$R = \{(1, 1), (1, 2), (1, 3), (3, 2)\}$$

then

$$R^{-1} = \{(1, 1), (2, 1), (3, 1), (2, 3)\}$$

5. **Complement of a relation** : Let relation  $R$  is defined from  $A$  to  $B$ , then complement  $R$  is set of ordered pairs  $\{(a, b) : (a, b) \notin R\}$ . It is also called complementary relation.

**For example** :

$$\text{Let } A = \{1, 2, 3\} \quad B = \{4, 5\}$$

$$\text{Then } A \times B = \{(1, 4), (1, 5), (2, 4), (2, 5), (3, 4), (3, 5)\}$$

$$\text{Let } R \text{ be defined as } R = \{(1, 4), (3, 4), (3, 5)\}$$

$$\text{Then } R^c = \bar{R} = \{(1, 5), (2, 4), (2, 5)\}$$

**Que 1.12.** Explain operation on relation with example.

**Answer**

1. Relations are sets of ordered pairs so all set operations can be done on relations.
2. The resulting sets contain ordered pairs and are, therefore, relations.
3. If  $R$  and  $S$  denote two relations, then  $R \cap S$ , known as intersection of  $R$  and  $S$ , defines a relation such that

$$x(R \cap S) y = x R y \wedge x S y$$

4. Similarly,  $R \cup S$ , known as union of  $R$  and  $S$ , such that

$$x(R \cup S) y = x R y \vee x S y$$

Also,  $x(R - S) y = x R y \wedge x \notin S y$  where  $R - S$  is known as different of  $R$  and  $S$

and  $x(R^c) y = x \notin R y$  where  $R^c$  is the complement of  $R$

**For example** :  $A = \{x, y, z\}$ ,  $B = \{x, y, z\}$ ,  $C = \{x, y, z\}$

$$D = \{Y, z\}, R = \{(x, X), (x, Y), (y, z)\}$$

$$S = \{(x, Y), (y, z)\}$$

The complement of  $R$  consists of all pairs of the Cartesian product  $A \times B$  that are not  $R$ . Thus  $A \times B = \{(x, X), (x, Y), (x, Z), (y, X), (y, Y), (y, Z), (z, X), (z, Y), (z, Z)\}$

Hence

$$R^c = \{(x, Z), (y, X), (y, Y), (z, X), (z, Y), (z, Z)\}$$

$$R \cup S = \{(x, X), (x, Y), (y, Z)\}$$

$$R \cap S = \{(x, Y), (y, Z)\}$$

$$R - S = \{(x, Y)\}$$

**Que 1.13.** Give properties of relation.

**Answer**

**Properties of relation are :**

1. **Reflexive relation** : A binary relation  $R$  on set  $A$  is said to be reflexive if every element of set  $A$  is related to itself.

*i.e.*,  $\forall a \in A, (a, a) \in R$  or  $aRa$ .

**For example :** Let  $R = \{(1, 1), (1, 2), (2, 2), (2, 3), (3, 3)\}$  be a relation defined on set  $A = \{1, 2, 3\}$ . As  $(1, 1) \in R, (2, 2) \in R$  and  $(3, 3) \in R$ . Therefore,  $R$  is reflexive relation.

2. **Irreflexive relation :** A binary relation  $R$  defined on set  $A$  is said to be irreflexive if there is no element in  $A$  which is related to itself *i.e.*,

$\forall a \in A$  such that  $(a, a) \notin R$ .

**For example :** Let  $R = \{(1, 2), (2, 1), (3, 1)\}$  be a relation defined on set  $A = \{1, 2, 3\}$ . As  $(1, 1) \notin R, (2, 2) \notin R$  and  $(3, 3) \notin R$ . Therefore,  $R$  is irreflexive relation.

3. **Non-reflexive relation :** A relation  $R$  defined on set  $A$  is said to be non-reflexive if it is neither reflexive nor irreflexive *i.e.*, some elements are related to itself but there exist at least one element not related to itself.

4. **Symmetric relation :** A binary relation on a set  $A$  is said to be symmetric if  $(a, b) \in R \Rightarrow (b, a) \in R$ .

5. **Asymmetric relation :** A binary relation on a set  $A$  is said to be asymmetric if  $(a, b) \in R \Rightarrow (b, a) \notin R$ .

6. **Antisymmetric relation :** A binary relation  $R$  defined on a set  $A$  is said to be antisymmetric relation if  $(a, b) \in R$  and  $(b, a) \in R \Rightarrow a = b$  *i.e.*,  $aRb$  and  $bRa \Rightarrow a = b$  for  $a, b \in R$ .

7. **Transitive relation :** A binary relation  $R$  on a set  $A$  is transitive whenever  $(a, b) \in R$  and  $(b, c) \in R$  then  $(a, c) \in R$   
*i.e.*,  $aRb$  and  $bRc \Rightarrow aRc$ .

**Que 1.14.** Write short notes on :

- Equivalence relation
- Composition of relation

OR

Write a short note on equality of relation.

**Answer**

- Equivalence relation :**

- A relation  $R$  on a set  $A$  is said to be equivalence relation if it is reflexive, symmetric and transitive.
- The two elements  $a$  and  $b$  related by an equivalence relation are called equivalent.
- So, a relation  $R$  is called equivalence relation on set  $A$  if it satisfies following three properties :

i.  $(a, a) \in R \forall a \in A$

(Reflexive)

ii.  $(a, b) \in R \Rightarrow (b, a) \in R$

(Symmetric)

iii.  $(a, b) \in R$  and  $(b, c) \in R \Rightarrow (a, c) \in R$

(Transitive)

- Composition of relation :**

- Let  $R$  be a relation from a set  $A$  to  $B$  and  $S$  be a relation from set  $B$  to  $C$  then composition of  $R$  and  $S$  is a relation consisting or ordered pair  $(a, c)$  where  $a \in A$  and  $c \in C$  provided that there exist  $b \in B$  such that  $(a, b) \in R \subseteq A \times B$  and  $(b, c) \in S \subseteq B \times C$ . It is denoted by  $R \circ S$ .

2. Symbolically,  $R \circ S = \{(a, c) \mid \exists b \in B \text{ such that } (a, b) \in R \text{ and } (b, c) \in S\}$

**Que 1.15.** Describe recursive definition of relation.

**Answer**

- The characteristic function  $C_R$  of a relation  $R \subseteq N^k$  is defined as follows :
  - $C_S(X_1, \dots, X_k) = 1$  if  $\langle X_1, \dots, X_k \rangle \in S$
  - $C_S(X_1, \dots, X_k) = 0$  if  $\langle X_1, \dots, X_k \rangle \notin S$
- A relation  $R$  is a recursive set iff its characteristic function  $C_R$  is a recursive function.
- Examples of recursive relations :  $<, >, \leq, =$ 
  - $c_<(x, y) = sg(y \div x)$
  - $c_>(x, y) = sg(x \div y)$
  - $c_{\leq}(x, y) = \overline{sg}(x + y)$
  - $c_{=} (x, y) = \overline{sg}(x \div y) \times c_{\leq}(x, y) = \overline{sg}(y + x)$
- Consider relation  $R(x, y, z)$  defined as follows :
  - $R(x, y, z)$  iff  $y \times z \leq x$
- We see that  $R$  is the result of substituting the recursive function  $\times$  into recursive relation  $\leq$ .
- Thus,  $R$  is recursive.
- (Technically,  $R$  is the result of substituting the functions  $f_1(x, y, z) = y \times z$  and  $f_2(x, y, z) = x$  into  $\leq$ , and we need to show that  $f_1(x, y, z) = y \times z$  and  $f_2(x, y, z) = x$  are recursive ... but that's trivial using the identity functions).

**Que 1.16.** Define the term partial order relation or partial ordering relation.

**Answer**

A binary relation  $R$  defined on set  $A$  is called Partial Order Relation (POR) if  $R$  satisfies following properties :

- $(a, a) \in R \forall a \in A$  (Reflexive)
- If  $(a, b) \in R$  and  $(b, a) \in R$ , then  $a = b$  (Antisymmetric)
- If  $(a, b) \in R$  and  $(b, c) \in R \Rightarrow (a, c) \in R$  where  $a, b, c \in A$  (Transitive)

A set  $A$  together with a partial order relation  $R$  is called partial order set or poset.

**Que 1.17.** Write short notes on :

- Closure of relations
- Total order
- Compatibility relation

**Answer****a. Closure of relations :**

i. **Reflexive closure :** Let  $R$  be a relation defined on set  $A$ . The  $R \cup I_A$  is called reflexive closure of  $R$ , where  $I_A = \{(a, a) | a \in A\}$  is diagonal or identity relation.

ii. **Symmetric closure :** Let  $R$  be a relation defined on set  $A$ . Then  $R \cup R^{-1}$  is called symmetric closure of  $R$ , where  $R^{-1}$  is inverse of  $R$  on  $A$ .

**b. Total order :** A binary relation  $R$  on a set  $A$  is said to be total order iff it is

i. Partial order

ii.  $(a, b) \in R$  or  $(b, a) \in R \quad \forall a, b \in A$

It is also called linear order.

**c. Compatibility relation :** A binary relation  $R$  defined on set  $A$  is said to be compatible relation if it is reflexive and symmetric. It is denoted by ' $\approx$ '.

**Que 1.18.** Show that  $R = \{(a, b) | a \equiv b \pmod{m}\}$  is an equivalence relation on  $Z$ . Show that if  $x_1 \approx y_1$  and  $x_2 \approx y_2$  then  $(x_1 + x_2) \approx (y_1 + y_2)$ .

AKTU 2014-15, Marks 05

**Answer**

$$R = \{(a, b) | a \equiv b \pmod{m}\}$$

For an equivalence relation it has to be reflexive, symmetric and transitive.

**Reflexive :** For reflexive  $\forall a \in Z$  we have  $(a, a) \in R$  i.e.,

$$a \equiv a \pmod{m}$$

$\Rightarrow a - a$  is divisible by  $m$  i.e., 0 is divisible by  $m$

Therefore  $aRa, \forall a \in Z$ , it is reflexive.

**Symmetric :** Let  $(a, b) \in R$  and we have

$$(a, b) \in R \text{ i.e., } a \equiv b \pmod{m}$$

$\Rightarrow a - b$  is divisible by  $m$

$\Rightarrow a - b = km, k$  is an integer

$\Rightarrow (b - a) = (-k)m$

$\Rightarrow (b - a) = pm, p$  is also an integer

$\Rightarrow b - a$  is also divisible by  $m$

$\Rightarrow b \equiv a \pmod{m} \Rightarrow (b, a) \in R$

It is symmetric.

**Transitive :** Let  $(a, b) \in R$  and  $(b, c) \in R$  then

$(a, b) \in R \Rightarrow a - b$  is divisible by  $m$

$\Rightarrow a - b = tm, t$  is an integer

...(1.18.1)

$(b, c) \in R \Rightarrow b - c$  is divisible by  $m$

$\Rightarrow b - c = sm, s$  is an integer

...(1.18.2)

From eq. (1.18.1) and (1.18.2)

$$a - b + b - c = (t + s)m$$

$$a - c = lm, l \text{ is also an integer}$$

$a - c$  is divisible by  $m$

$a \equiv c \pmod{m}$ , yes it is transitive.

$R$  is an equivalence relation.

**To show :**  $(x_1 + x_2) \equiv (y_1 + y_2) :$

It is given  $x_1 \equiv y_1$  and  $x_2 \equiv y_2$

i.e.,  $x_1 - y_1$  divisible by  $m$

$x_2 - y_2$  divisible by  $m$

Adding above equation :

$(x_1 - y_1) + (x_2 - y_2)$  is divisible by  $m$

$\Rightarrow (x_1 + x_2) - (y_1 + y_2)$  is divisible by  $m$

i.e.,  $(x_1 + x_2) \equiv (y_1 + y_2)$

**Que 1.19.** Let  $R$  be binary relation on the set of all strings of 0's and 1's such that  $R = \{(a, b) \mid a \text{ and } b \text{ are strings that have the same number of 0's}\}$ . Is  $R$  is an equivalence relation and a partial ordering relation ?

AKTU 2014-15, Marks 05

**Answer**

**For equivalence relation :**

**Reflexive :**  $a R a \Rightarrow (a, a) \in R \forall a \in R$

where  $a$  is a string of 0's and 1's.

Always  $a$  is related to  $a$  because both  $a$  has same number of 0's.

It is reflexive.

**Symmetric :** Let  $(a, b) \in R$

then  $a$  and  $b$  both have same number of 0's which indicates that again both  $b$  and  $a$  will also have same number of zeros. Hence  $(b, a) \in R$ . It is symmetric.

**Transitive :** Let  $(a, b) \in R, (b, c) \in R$

$(a, b) \in R \Rightarrow a$  and  $b$  have same number of zeros.

$(b, c) \in R \Rightarrow b$  and  $c$  have same number of zeros.

Therefore  $a$  and  $c$  also have same number of zeros, hence  $(a, c) \in R$ .

It is transitive.

$\therefore R$  is an equivalence relation.

For partial order, it has to be reflexive, antisymmetric and transitive. Since, symmetricity and antisymmetricity cannot hold together. Therefore, it is not partial order relation.

**Que 1.20.** Let  $A = \{1, 2, 3, \dots, 13\}$ . Consider the equivalence relation on  $A \times A$  defined by  $(a, b) R (c, d)$  if  $a + d = b + c$ . Find equivalence classes of  $(5, 8)$ .

AKTU 2014-15, Marks 05

**Answer**

$$A = \{1, 2, 3, \dots, 13\}$$

$$[(5, 8)] = \{(a, b) : (a, b) R (5, 8), (a, b) \in A \times A\}$$

$$\begin{aligned}
 &= \{(a, b) : a + 8 = b + 5\} \\
 &= \{(a, b) : a + 3 = b\} \\
 [5, 8] &= \{(1, 4), (2, 5), (3, 6), (4, 7) \\
 &\quad (5, 8), (6, 9), (7, 10), (8, 11) \\
 &\quad (9, 12), (10, 13)\}
 \end{aligned}$$

**Que 1.21.** The following relation on  $A = \{1, 2, 3, 4\}$ . Determine whether the following :

- a.  $R = \{(1, 3), (3, 1), (1, 1), (1, 2), (3, 3), (4, 4)\}$   
 b.  $R = A \times A$

Is an equivalence relation or not ?

**AKTU 2015-16, Marks 10**

**Answer**

- a.  $R = \{(1, 3), (3, 1), (1, 1), (1, 2), (3, 3), (4, 4)\}$

**Reflexive :**  $(a, a) \in R \forall a \in A$

$$\because (1, 1) \in R, (2, 2) \notin R$$

$\therefore R$  is not reflexive.

**Symmetric :** Let  $(a, b) \in R$  then  $(b, a) \in R$ .

$$\because (1, 3) \in R \text{ so } (3, 1) \in R$$

$$\because (1, 2) \in R \text{ but } (2, 1) \notin R$$

$\therefore R$  is not symmetric.

**Transitive :** Let  $(a, b) \in R$  and  $(b, c) \in R$  then  $(a, c) \in R$

$$\because (1, 3) \in R \text{ and } (3, 1) \in R \text{ so } (1, 1) \in R$$

$$\because (2, 1) \in R \text{ and } (1, 3) \in R \text{ but } (2, 3) \notin R$$

$\therefore R$  is not transitive.

Since,  $R$  is not reflexive, not symmetric, and not transitive so  $R$  is not an equivalence relation.

- b.  $R = A \times A$

Since,  $A \times A$  contains all possible elements of set  $A$ . So,  $R$  is reflexive, symmetric and transitive. Hence  $R$  is an equivalence relation.

**Que 1.22.** Let  $(A, \leq)$  be a partially ordered set. Let  $\leq$  be a binary relation  $A$  such that for  $a$  and  $b$  in  $A$ ,  $a$  is related to  $b$  iff  $b \leq a$ .

- i. Show that  $\leq$  is partially ordered relation.  
 ii. Show that  $(A, \leq)$  is lattice or not.

**Answer**

- i.  $(A, \leq)$  is a partially ordered relation if it is reflexive, antisymmetric and transitive.

**Reflexive :** Let  $a \in A$  then by definition of relation,  $aRa \Rightarrow a \leq a$  which is true.

Hence, the relation  $R$  i.e.,  $\leq$  is reflexive.

**Antisymmetric :** Let  $a, b \in A$  and

$$aRb \Rightarrow b \leq a$$

$\Rightarrow$

$$a \leq b$$

$$\Rightarrow b R a$$

$aRb$  and  $bRa$  holds only when  $a = b$ . Relation is antisymmetric.

**Transitive :** Let  $a, b, c \in A$  and  $aRb$  and  $bRc$

$$\Rightarrow b \leq a, c \leq b$$

$$\Rightarrow c \leq a$$

$$\Rightarrow aRc$$

Hence, relation is transitive.

Therefore,  $\leq$  is a partial order relation.

- ii. Since all the elements of the given set  $A$  are comparable to each other, we always have a least upper bound and greatest lower bound for each pair of elements of  $A$  and  $A$  is also a partial order relation. Hence,  $(A, \leq)$  is a lattice.

**Que 1.23.** Let  $n$  be a positive integer and  $S$  a set of strings. Suppose that  $R_n$  is the relation on  $S$  such that  $sR_n t$  if and only if  $s = t$ , or both  $s$  and  $t$  have at least  $n$  characters and first  $n$  characters of  $s$  and  $t$  are the same. That is, a string of fewer than  $n$  characters is related only to itself; a string  $s$  with at least  $n$  characters is related to a string  $t$  if and only if  $t$  has at least  $n$  characters and  $t$  begins with the  $n$  characters at the start of  $s$ .

**AKTU 2018-19, Marks 07**

**Answer**

We have to show that the relation  $R_n$  is reflexive, symmetric, and transitive.

- 1. Reflexive :** The relation  $R_n$  is reflexive because  $s = s$ , so that  $sR_n s$  whenever  $s$  is a string in  $S$ .
- 2. Symmetric :** If  $sR_n t$ , then either  $s = t$  or  $s$  and  $t$  are both at least  $n$  characters long that begin with the same  $n$  characters. This means that  $tR_n s$ . We conclude that  $R_n$  is symmetric.
- 3. Transitive :** Now suppose that  $sR_n t$  and  $tR_n u$ . Then either  $s = t$  or  $s$  and  $t$  are at least  $n$  characters long and  $s$  and  $t$  begin with the same  $n$  characters, and either  $t = u$  or  $t$  and  $u$  are at least  $n$  characters long and  $t$  and  $u$  begin with the same  $n$  characters. From this, we can deduce that either  $s = u$  or both  $s$  and  $u$  are  $n$  characters long and  $s$  and  $u$  begin with the same  $n$  characters, i.e.,  $sR_n u$ . Consequently,  $R_n$  is transitive.

**Que 1.24.** Let  $X = \{1, 2, 3, \dots, 7\}$  and  $R = \{(x, y) \mid (x - y) \text{ is divisible by } 3\}$ .

Is  $R$  equivalence relation. Draw the digraph of  $R$ .

**AKTU 2017-18, Marks 07**

**Answer**

Given that

$$X = \{1, 2, 3, 4, 5, 6, 7\}$$

and

$$R = \{(x, y) : (x - y) \text{ is divisible by } 3\}$$

Then  $R$  is an equivalence relation if

i. **Reflexive:**  $\forall x \in X \Rightarrow (x - x)$  is divisible by 3

So,  $(x, x) \in R \forall x \in X$  or,  $R$  is reflexive.

ii. **Symmetric:** Let  $x, y \in X$  and  $(x, y) \in R$

$\Rightarrow (x - y)$  is divisible by 3  $\Rightarrow (x - y) = 3n_1$ , ( $n_1$  being an integer)

$\Rightarrow (y - x) = -3n_1 = 3n_2$ ,  $n_2$  is also an integer

So,  $y - x$  is divisible by 3 or  $R$  is symmetric.

iii. **Transitive:** Let  $x, y, z \in X$  and  $(x, y) \in R$ ,  $(y, z) \in R$

Then  $x - y = 3n_1$ ,  $y - z = 3n_2$ ,  $n_1, n_2$  being integers

$\Rightarrow x - z = 3(n_1 + n_2)$ ,  $n_1 + n_2 = n_3$  be any integer

So,  $(x - z)$  is also divisible by 3 or  $(x, z) \in R$

So,  $R$  is transitive.

Hence,  $R$  is an equivalence relation.




Fig. 1.24.1. Diagram of  $R$ .

### PART-4

*Function : Definition, Classification of Functions, Operations on Functions, Recursively Defined Functions, Growth of Functions.*

#### Questions-Answers

Long Answer Type and Medium Answer Type Questions

**Que 1.25.** Define the term function. Also, give classification of it.

#### Answer

- Let  $X$  and  $Y$  be any two non-empty sets. A function from  $X$  to  $Y$  is a rule that assigns to each element  $x \in X$  a unique element  $y \in Y$ .
- If  $f$  is a function from  $X$  to  $Y$  we write  $f: X \rightarrow Y$ .
- Functions are denoted by  $f, g, h, i$  etc.
- It is also called mapping or transformation or correspondence.

**Domain and co-domain of a function:** Let  $f$  be a function from  $X$  to  $Y$ . Then set  $X$  is called domain of function  $f$  and  $Y$  is called co-domain of function  $f$ .

**Range of function :** The range of  $f$  is set of all images of elements of  $X$ .

i.e.,  $\text{Range}(f) = \{y : y \in Y \text{ and } y = f(x) \forall x \in X\}$

Also  $\text{Range}(f) \subseteq Y$

**Classification of functions :**

1. **Algebraic functions :** Algebraic functions are those functions which consist of a finite number of terms involving powers and roots of the independent variable  $x$ .

Three particular cases of algebraic functions are :

i. **Polynomial functions :** A function of the form  $a_0x^n + a_1x^{n-1} + \dots + a_n$  where  $n$  is a positive integer and  $a_0, a_1, \dots, a_n$  are real constants and  $a_0 \neq 0$  is called a polynomial of  $x$  in degree  $n$ , for example  $f(x) = 2x^3 + 5x^2 + 7x - 3$  is a polynomial of degree 3.

ii. **Rational functions :** A function of the form  $\frac{f(x)}{g(x)}$  where  $f(x)$  and  $g(x)$  are polynomials.

iii. **Irrational functions :** Functions involving radicals are called irrational functions.  $f(x) = \sqrt[3]{x} + 5$  is an example of irrational function.

2. **Transcendental functions :** A function which is not algebraic is called transcendental function.

i. **Trigonometric functions :** Six functions  $\sin x, \cos x, \tan x, \sec x, \text{cosec } x, \cot x$  where the angle  $x$  is measured in radian are called trigonometric functions.

ii. **Inverse trigonometric functions :** Six functions  $\sin^{-1} x, \cos^{-1} x, \tan^{-1} x, \cot^{-1} x, \sec^{-1} x, \text{cosec}^{-1} x$ , are called inverse trigonometric functions.

iii. **Exponential functions :** A function  $f(x) = a^x$  ( $a > 0$ ) satisfying the law  $a^x = a$  and  $a^x a^y = a^{x+y}$  is called the exponential function.


iv. **Logarithm functions :** The inverse of the exponential function is called logarithm function.

**Que 1.26.** Give the types / operations on functions.

**Answer**

1. **One-to-one function (Injective function or injection) :** Let  $f: X \rightarrow Y$  then  $f$  is called one-to-one function if for distinct elements of  $X$  there are distinct image in  $Y$  i.e.,  $f$  is one-to-one iff

$$f(x_1) = f(x_2) \text{ implies } x_1 = x_2 \forall x_1, x_2 \in X$$



**Fig. 1.26.1. One-to-one.**

2. **Onto function (Surjection or surjective function)** : Let  $f: X \rightarrow Y$  then  $f$  is called onto function iff for every element  $y \in Y$  there is an element  $x \in X$  with  $f(x) = y$  or  $f$  is onto if  $\text{Range}(f) = Y$ .




Fig. 1.26.2. Onto.

3. **One-to-one onto function (Bijective function or bijection)** : A function which is both one-to-one and onto is called one-to-one onto function or bijective function.




Fig. 1.26.3. One-to-one onto.

4. **Many one function** : A function which is not one-to-one is called many one function i.e., two or more elements in domain have same image in co-domain i.e.,

$$\text{If } f: X \rightarrow Y \text{ then } f(x_1) = f(x_2) \Rightarrow x_1 \neq x_2.$$




Fig. 1.26.4. Many one.

5. **Identity function** : Let  $f: X \rightarrow X$  then  $f$  is called identity function if  $f(a) = a \forall a \in X$  i.e., every element of  $X$  is image of itself. It is denoted by  $I$ .
6. **Inverse function (Invertible function)** : Let  $f$  be a bijective function from  $X$  to  $Y$ . The inverse function of  $f$  is the function that assigns an element  $y \in Y$ , a unique element  $x \in X$  such that  $f(x) = y$  and inverse of  $f$  denoted by  $f^{-1}$ . Therefore if  $f(x) = y$  implies  $f^{-1}(y) = x$ .

**Que 1.27.** Determine whether each of these functions is a bijective from  $R$  to  $R$ .

a.  $f(x) = x^2 + 1$

b.  $f(x) = x^3$

c.  $f(x) = (x^2 + 1)/(x^2 + 2)$

AKTU 2015-16, Marks 15

**Answer**

a.  $f(x) = x^2 + 1$

Let  $x_1, x_2 \in R$  such that

$$\begin{aligned} f(x_1) &= f(x_2) \\ x_1^2 + 1 &= x_2^2 + 1 \\ x_1^2 &= x_2^2 \\ x_1 &= \pm x_2 \end{aligned}$$

Therefore, if  $x_2 = 1$  then  $x_1 = \pm 1$

So,  $f$  is not one-to-one.

Hence,  $f$  is not bijective.

b. Let  $x_1, x_2 \in R$  such that  $f(x_1) = f(x_2)$

$$\begin{aligned} x_1^3 &= x_2^3 \\ x_1 &= x_2 \end{aligned}$$

$\therefore f$  is one-to-one.

Let  $y \in R$  such that

$$\begin{aligned} y &= x^3 \\ x &= (y)^{1/3} \end{aligned}$$

For  $\forall y \in R \exists$  a unique  $x \in R$  such that  $y = f(x)$

$\therefore f$  is onto.

Hence,  $f$  is bijective.

c. Let  $x_1, x_2 \in R$  such that  $f(x_1) = f(x_2)$

$$\Rightarrow \frac{x_1^2 + 1}{x_1^2 + 2} = \frac{x_2^2 + 1}{x_2^2 + 2}$$

If  $x_1 = 1, x_2 = -1$  then  $f(x_1) = f(x_2)$   
but  $x_1 \neq x_2$

$\therefore f$  is not one-to-one.

Hence,  $f$  is not bijective.

**Que 1.28.** If  $f: A \rightarrow B, g: B \rightarrow C$  are invertible functions, then show that  $g \circ f: A \rightarrow C$  is invertible and  $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$ .

**AKTU 2014-15, Marks 05**

**Answer**

If  $f: A \rightarrow B$  and  $g: B \rightarrow C$  be one-to-one onto functions, then  $g \circ f$  is also one-to-one and  $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$

**Proof.** Since  $f$  is one-to-one,  $f(x_1) = f(x_2) \Rightarrow x_1 = x_2$  for  $x_1, x_2 \in R$

Again since  $g$  is one-to-one,  $g(y_1) = g(y_2) \Rightarrow y_1 = y_2$  for  $y_1, y_2 \in R$

Now  $g \circ f$  is one-to-one, since  $(g \circ f)(x_1) = (g \circ f)(x_2) \Rightarrow g[f(x_1)] = g[f(x_2)]$

$$\Rightarrow f(x_1) = f(x_2) \quad [g \text{ is one-to-one}]$$

$$\Rightarrow x_1 = x_2 \quad [f \text{ is one-to-one}]$$

Since  $g$  is onto, for  $z \in C$ , there exists  $y \in B$  such that  $g(y) = z$ . Also  $f$  being onto there exists  $x \in A$  such that  $f(x) = y$ . Hence  $z = g(y)$

$$= g[f(x)] = (g \circ f)(x)$$

This shows onto.

Thus,  $g \circ f$

By the de

$(g \circ f)^{-1}: C$

Also  $g^{-1}: C$

Then by th

Therefore,

Now

Thus,

So,

**Que 1.29**

a. Com

b. Rect

c. Prin

**Answer**

a. Com

If:

from

1.

2.

3.

b. Re

So

be

as

This shows that every element  $z \in C$  has pre-image under  $g \circ f$ . So,  $g \circ f$  is onto.

Thus,  $g \circ f$  is one-to-one onto function and hence  $(g \circ f)^{-1}$  exists.

By the definition of the composite functions,  $g \circ f : A \rightarrow C$ . So,  $(g \circ f)^{-1} : C \rightarrow A$ .

Also  $g^{-1} : C \rightarrow B$  and  $f^{-1} : B \rightarrow A$ .

Then by the definition of composite functions,  $f^{-1} \circ g^{-1} : C \rightarrow A$ .

Therefore, the domain of  $(g \circ f)^{-1}$  = the domain of  $f^{-1} \circ g^{-1}$ .

$$\begin{aligned} \text{Now } (g \circ f)^{-1}(z) = x &\Leftrightarrow (g \circ f)(x) = z \\ &\Leftrightarrow g(f(x)) = z \\ &\Leftrightarrow g(y) = z \text{ where } y = f(x) \\ &\Leftrightarrow y = g^{-1}(z) \\ &\Leftrightarrow f^{-1}(y) = f^{-1}(g^{-1}(z)) = (f^{-1} \circ g^{-1})(z) \\ &\Leftrightarrow x = (f^{-1} \circ g^{-1})(z) \quad [f^{-1}(y) = x] \end{aligned}$$

Thus,  $(g \circ f)^{-1}(z) = (f^{-1} \circ g^{-1})(z)$ .

So,  $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$ .


**Que 1.29.** Explain the following :

- Composition of functions
- Recursive function
- Primitive recursion

**Answer**

- Composition of functions :**

If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  then the composition of  $f$  and  $g$  is a new function from  $X$  to  $Z$  denoted by  $g \circ f = g(f(x))$



**Fig. 1.29.1.** Many one onto.

- If  $f$  and  $g$  are considered to be relations then composition of  $f$  and  $g$  was denoted by  $f \circ g$  whereas the composition of function  $f$  and  $g$  is denoted by  $g \circ f$ .
  - Composition of functions is not commutative i.e.,  $f \circ g \neq g \circ f$ .
  - Composition of functions is associative i.e.,  $f \circ (g \circ h) = (f \circ g) \circ h$ , where  $h : X \rightarrow Y$ ,  $g : Y \rightarrow Z$ , and  $f : Z \rightarrow W$ .
- Recursive function (Recursively defined function) :**  
Sometimes it becomes difficult to define a function explicitly then it may be easy to define this function in terms of itself. This technique is referred as recursion.

This concept of recursion can be used to define sets, sequences and algorithms also.

We will define three functions which are called Basis function or Initial functions that are used in defining other functions by induction.

1. **Zero function** : ( $Z : z(x) = 0$ )
2. **Successor function** : ( $S : s(x) = x + 1$ )
3. **Projection function or generalized identity function** :

$$U_i^n : U_i^n(x_1, x_2, \dots, x_n) = x_i$$

where superscripts  $n$  indicates a number of argument of the function and subscripts  $i$  indicates the value of the function is equal to  $i^{\text{th}}$  argument.

**c. Primitive recursion :**

A function is called primitive recursion if and only if it can be constructed from the initial functions and other known primitive recursive functions by a finite number of operations and recursion only.

**For example :** Consider the function

$$\begin{aligned} f(x, y) &= x + y \\ \text{Now } f(x, y + 1) &= x + (y + 1) \\ &= (x + y) + 1 \\ &= f(x, y) + 1 \\ \text{Also } f(x, 0) &= x + 0 = x \\ \text{Now } f(x, 0) &= x = U_1^1(x) \quad \dots(1.29.1) \\ \text{Also } f(x, y + 1) &= f(x, y) + 1 \end{aligned}$$

$$\begin{aligned} &= S(f(x, y)) \text{ where } S \text{ is the successor function} \\ &= (U_3^3[x, y, f(x, y)]) \quad \dots(1.29.2) \end{aligned}$$

If we consider  $g(x) = U_1^1(x)$  and  $h(x, y, z) = S \cdot U_3^3(x, y, z)$

From eq. (1.29.1) and (1.29.2), we get  $f(x, 0) = g(x)$

and  $f(x, y + 1) = h(x, y, f(x, y))$

Thus,  $f$  is obtained from the initial functions  $U_1^1$ ,  $U_3^3$  and  $S$  by applying compositions once and recursion once. Hence,  $f$  is primitive recursive.

**Que 1.30.** Write short note on growth of functions.

**Answer**

1. We need to approximate the number of steps required to execute any algorithm because of the difficulty involved in expression or difficulty in evaluating an expression. We compare one function with another function to know their rate of growths.
2. If  $f$  and  $g$  are two functions we can give the statements like ' $f$  has same growth rate as  $g$ ' or ' $f$  has higher growth rate than  $g$ '.
3. Growth rate of function can be defined through notation.
  - a.  **$\theta$ -Notation (Same order)** : This notation bounds a function to within constant factors. We say  $f(n) = \theta g(n)$  if there exist positive constants  $n_0$ ,  $c_1$  and  $c_2$  such that to the right of  $n_0$  the value of  $f(n)$  always lies between  $c_1 g(n)$  and  $c_2 g(n)$  inclusive.




Fig. 1.30.1.

- b. **O-Notation (Upper bound)** : This notation gives an upper bound for a function to within a constant factor. We write  $f(n) = O(g(n))$  if there are positive constants  $n_0$  and  $c$  such that to the right of  $n_0$ , the value of  $f(n)$  always lies on or below  $cg(n)$ .




Fig. 1.30.2.

- c. **Ω-Notation (Lower bound)** : This notation gives a lower bound for a function to within a constant factor. We write  $f(n) = Ωg(n)$  if there are positive constants  $n_0$  and  $c$  such that to the right of  $n_0$ , the value of  $f(n)$  always lies on or above  $cg(n)$ .




Fig. 1.30.3.

### PART-5

*Natural Numbers : Introduction, Mathematical Induction, Variants of Induction, Induction with Non-zero Base Cases.*

## Questions-Answers

## Long Answer Type and Medium Answer Type Questions

**Que 1.31.** Describe mathematical induction.

## Answer

Mathematical induction is a technique of proving a proposition over the positive integers. It is the most basic method of proof used for proving statements having a general pattern. A formal statement of principle of mathematical induction can be stated as follows :

Let  $S(n)$  be statement that involve positive integer  $n = 1, 2, \dots$  then

**Step 1 :** Verify  $S(1)$  is true. (Inductive base)

**Step 2 :** Assume that  $S(k)$  is true for some arbitrary  $k$ . (Inductive hypothesis)

**Step 3 :** Verify  $S(k + 1)$  is true using basis of inductive hypothesis. (Inductive step)

**Que 1.32.** Prove that  $n^3 + 2n$  is divisible by 3 using principle of mathematical induction, where  $n$  is natural number.

AKTU 2015-16, Marks 10

## Answer

Let  $S(n) : n^3 + 2n$  is divisible by 3.

**Step I : Inductive base :** For  $n = 1$

$$(1)^3 + 2 \cdot 1 = 3 \text{ which is divisible by 3}$$

Thus,  $S(1)$  is true.

**Step II : Inductive hypothesis :** Let  $S(k)$  is true i.e.,  $k^3 + 2k$  is divisible by 3 holds true.

$$\text{or } k^3 + 2k = 3s \text{ for } s \in N$$

**Step III : Inductive step :** We have to show that  $S(k + 1)$  is true

i.e.,  $(k + 1)^3 + 2(k + 1)$  is divisible by 3

Consider  $(k + 1)^3 + 2(k + 1)$

$$= k^3 + 1 + 3k^2 + 3k + 2k + 2$$

$$= (k^3 + 2k) + 3(k^2 + k + 1)$$

$$= 3s + 3l \text{ where } l = k^2 + k + 1 \in N$$

$$= 3(s + l)$$

Therefore,  $S(k + 1)$  is true

Hence by principle of mathematical induction  $S(n)$  is true for all  $n \in N$ .

**Que 1.33.** Prove by induction :  $\frac{1}{1.2} + \frac{1}{2.3} + \dots + \frac{1}{n(n+1)} = \frac{n}{(n+1)}$ .

**AKTU 2016-17, Marks 10**

**Answer**

Let the given statement be denoted by  $S(n)$ .

1. **Inductive base :** For  $n = 1$

$$\frac{1}{1.2} = \frac{1}{1+1} = \frac{1}{2}$$

Hence  $S(1)$  is true.

2. **Inductive hypothesis :** Assume that  $S(k)$  is true i.e.,

$$\frac{1}{1.2} + \frac{1}{2.3} + \dots + \frac{1}{k(k+1)} = \frac{k}{k+1}$$

3. **Inductive step :** We wish to show that the statement is true for  $n = k + 1$  i.e.,

$$\frac{1}{1.2} + \frac{1}{2.3} + \dots + \frac{1}{(k+1)(k+2)} = \frac{k+1}{k+2}$$

$$\text{Now, } \frac{1}{1.2} + \frac{1}{2.3} + \dots + \frac{1}{k(k+1)} + \frac{1}{(k+1)(k+2)}$$

$$= \frac{k}{k+1} + \frac{1}{(k+1)(k+2)} = \frac{k^2 + 2k + 1}{(k+1)(k+2)}$$

$$= \frac{k+1}{k+2}$$

Thus,  $S(k+1)$  is true whenever  $S(k)$  is true. By principle of mathematical induction,  $S(n)$  is true for all positive integer  $n$ .

**Que 1.34.** Prove by the principle of mathematical induction, that the sum of finite number of terms of a geometric progression,  $a + ar + ar^2 + \dots + ar^{n-1} = a(r^n - 1)/(r - 1)$  if  $r \neq 1$ .

**AKTU 2014-15, Marks 05**

**Answer**

**Basis :** True for  $n = 1$  i.e.,

$$\text{L.H.S} = a$$

$$\text{R.H.S} = \frac{a(r-1)}{r-1} = a$$

Therefore, L.H.S. = R.H.S.

**Induction :** Let it be true for  $n = k$  i.e.,

$$a + ar + ar^2 + \dots + ar^{k-1} = \frac{a(r^k - 1)}{r - 1} \quad \dots(1.34.1)$$

Now we will show that it is true for  $n = k + 1$  using eq. (1.34.1)

i.e.,  $a + ar + ar^2 + \dots + ar^{k-1} + ar^k$

Using eq. (1.34.1), we get

$$\begin{aligned} & \frac{a(r^k - 1)}{r - 1} + ar^k \\ &= \frac{ar^k - a + ar^{k+1} - ar^k}{r - 1} = \frac{a(r^{k+1} - 1)}{r - 1} \end{aligned}$$

which is R.H.S. for  $n = k + 1$ , hence it is true for  $n = k + 1$ .

By mathematical induction, it is true for all  $n$ .

**Que 1.35.** Prove that if  $n$  is a positive integer, then 133 divides

$$11^{n+1} + 12^{2n-1}.$$

**AKTU 2018-19, Marks 07**

**Answer**

We prove this by induction on  $n$ .

**Base case :** For  $n = 1$ ,  $11^{n+1} + 12^{2n-1} = 11^2 + 12^1 = 133$  which is divisible by 133.

**Inductive step :** Assume that the hypothesis holds for  $n = k$ , i.e.,  $11^{k+1} + 12^{2k-1} = 133A$  for some integer  $A$ . Then for  $n = k + 1$ ,

$$\begin{aligned} 11^{n+1} + 12^{2n-1} &= 11^{k+1+1} + 12^{2(k+1)-1} \\ &= 11^{k+2} + 12^{2k+1} \\ &= 11 * 11^{k+1} + 144 * 12^{2k-1} \\ &= 11 * 11^{k+1} + 11 * 12^{2k-1} + 133 * 12^{2k-1} \\ &= 11[11^{k+1} + 12^{2k-1}] + 133 * 12^{2k-1} \\ &= 11 * 133A + 133 * 12^{2k-1} \\ &= 133[11A + 12^{2k-1}] \end{aligned}$$

Thus if the hypothesis holds for  $n = k$  it also holds for  $n = k + 1$ . Therefore, the statement given in the equation is true.

**Que 1.36.** Prove by mathematical induction

$n^4 - 4n^2$  is divisible by 3 for all  $n > = 2$ .

**AKTU 2017-18, Marks 07**

**Answer**

**Base case :** If  $n = 0$ , then  $n^4 - 4n^2 = 0$ , which is divisible by 3.

**Inductive hypothesis :** For some  $n \geq 0$ ,  $n^4 - 4n^2$  is divisible by 3.

**Inductive step :** Assume the inductive hypothesis is true for  $n$ . We need to show that  $(n + 1)^4 - 4(n + 1)^2$  is divisible by 3. By the inductive hypothesis, we know that  $n^4 - 4n^2$  is divisible by 3.

Hence  $(n + 1)^4 - 4(n + 1)^2$  is divisible by 3 if  $(n + 1)^4 - 4(n + 1)^2 - (n^4 - 4n^2)$  is divisible by 3.

Now  $(n + 1)^4 = n^4 + 4n^3 + 6n^2 + 4n + 1$   
 $= 4n^3 + 6n^2 + 4n + 1$   
 which is divisible by 3 for all  $n \in \mathbb{N}$ .  
 Going back to the inductive hypothesis, we see that the inductive hypothesis holds for  $n + 1$ .  
 By the Principle of Mathematical Induction,  $n \in \mathbb{N}$ .

**Que 1.37.**

$$3 + 33 + 333$$

**Answer**

$$3 + 33 + 333$$

Let given

1. Indu

2. Indu

$$3 + 33$$

3. Indu

$$3 + 33$$

Now

$$= 3 +$$

$$= (10$$

$$= (10$$

$$= (10$$

Thu

mat

**Proof**

**Que 1**

$$\begin{aligned} & \text{Now } (n + 1)^4 - 4(n + 1)^2 - (n^4 - 4n^2) \\ &= n^4 + 4n^3 + 6n^2 + 4n + 1 - 4n^2 - 8n - 4 - n^4 + 4n^2 \\ &= 4n^3 + 6n^2 - 4n - 3, \end{aligned}$$

which is divisible by 3 if  $4n^3 - 4n$  is. Since  $4n^3 - 4n = 4n(n + 1)(n - 1)$ , we see that  $4n^3 - 4n$  is always divisible by 3.

Going backwards, we conclude that  $(n + 1)^4 - 4(n + 1)^2$  is divisible by 3, and that the inductive hypothesis holds for  $n + 1$ .

By the Principle of Mathematical Induction,  $n^4 - 4n^2$  is divisible by 3, for all  $n \in \mathbb{N}$ .

**Que 1.37.** Prove by mathematical induction

$$3 + 33 + 333 + \dots + 3333 = (10^{n+1} - 9n - 10)/27$$

AKTU 2017-18, Marks 07

Answer

$$3 + 33 + 333 + \dots + 3333 \dots = (10^{n+1} - 9n - 10)/27$$

Let given statement be denoted by  $S(n)$

1. **Inductive base :** For  $n = 1$

$$3 = \frac{(10^2 - 9(1) - 10)}{27}, 3 = \frac{100 - 19}{27} = \frac{81}{27} = 3$$

$3 = 3$ . Hence  $S(1)$  is true.

2. **Inductive hypothesis :** Assume that  $S(k)$  is true i.e.,

$$3 + 33 + 333 + \dots + 3333 = (10^{k+1} - 9k - 10)/27$$

3. **Inductive steps :** We have to show that  $S(k + 1)$  is also true i.e.,

$$3 + 33 + 333 + \dots + (10^{k+2} - 9(k+1) - 10)/27$$

$$\text{Now, } 3 + 33 + \dots + 33 \dots 3$$

$$= 3 + 33 + 333 + \dots + 3 \dots 3$$

$$= (10^{k+1} - 9k - 10)/27 + 3(10^{k+1} - 1)/9$$

$$= (10^{k+1} + 9k - 10 + 9 \cdot 10^{k+1} - 9)/27$$

$$= (10^{k+1} + 9 \cdot 10^{k+1} - 9k - 8 - 10)/27 = (10^{k+2} - 9(k+1) - 10)/27$$

Thus  $S(k + 1)$  is true whenever  $S(k)$  is true. By the principle of mathematical induction  $S(n)$  true for all positive integer  $n$ .

**PART-6**  
*Proof Methods, Proof by Counter - Example, Proof by Contradiction.*

**Questions-Answers**  
**Long Answer Type and Medium Answer Type Questions**

**Que 1.38.** What are the different proof methods ?

**Answer**

**Different methods of proof are :**

1. **Direct proof :** In this method, we will assume that hypothesis  $p$  is true from the implication  $p \rightarrow q$ . We can proof that  $q$  is true by using rules of interference or some other theorem. This will show  $p \rightarrow q$  is true. The combination of  $p$  is true and  $q$  is false will never occur.
2. **Indirect methods :** These are of five types as follows :
  - a. Proof by contrapositive.
  - b. Proof by contradiction (Reductio ad absurdum).
  - c. Proof by exhaustive cases.
  - d. Proof by cases.
  - e. Proof by counter example.

**Que 1.39.** Explain proof by counter example.

**Answer**

In this method, we will give an example for which given statement is false. This example is called counter example.

**For example :** Prove by counter that every positive integer can be written as sum of square of three integers.

To look for a counter example we will first write successive positive integers as the sum of square of three integers.

$$1 = 1^2 + 0^2 + 0^2$$

$$2 = 1^2 + 0^2 + 1^2$$

$$3 = 1^2 + 1^2 + 1^2$$

$$4 = 2^2 + 0^2 + 0^2$$

$$5 = 0^2 + 1^2 + 2^2$$

$$6 = 1^2 + 1^2 + 2^2$$

But there is no way to write 7 as a sum of three squares which gives counter example. Therefore, given statement is false.

**Que 1.40.** Explain proof by contradiction with example.

**Answer**

In this method, we assume that  $q$  is false i.e.,  $\sim q$  is true. Then by using rules of inference and other theorems we will show the given statement is true as well as false i.e., we will reach at a contradiction. Therefore,  $q$  must be true.

**For example :** Prove that  $\sqrt{3}$  is irrational.

Let  $\sqrt{3}$  is a rational number. Then  $\exists$  positive prime integer  $p$  and  $q$  such that

$$\sqrt{3} = \frac{p}{q}$$

$$\Rightarrow 3 = \frac{p^2}{q^2} \Rightarrow p^2 = 3q^2 \quad \dots(1.40.1)$$

$$\begin{aligned} &\Rightarrow 3/p^2 \text{ (3 divides } p^2) \\ &\Rightarrow 3/p \quad (\because p \text{ is an integer}) \\ \therefore &p = 3x \text{ for some } x \in Z \\ &\Rightarrow p^2 = 9x^2 \quad \dots(1.40.2) \end{aligned}$$

From eq. (1.40.1) and (1.40.2) we get,

$$\begin{aligned} &9x^2 = 3q^2 \\ &\Rightarrow q^2 = 3x^2 \\ &\Rightarrow 3/q^2 \text{ (3 divides } q^2) \\ &\Rightarrow 3/q \end{aligned}$$

$\therefore$  3 divides  $p$  and  $q$  which is contradiction to our assumption that  $p$  and  $q$  are prime. Therefore,  $\sqrt{3}$  is irrational.

**Que 1.41.** Write a short note on the following with example :

- i. Proof by contrapositive.
- ii. Proof by exhaustive cases.
- iii. Proof by cases.

**Answer**

- i. **Proof by contrapositive** : In this we can prove  $p \rightarrow q$  is true by showing  $\sim q \rightarrow \sim p$  is true. It is also called proof by contraposition.

**Example** : Using method of contraposition if  $n$  is integer and  $3n + 2$  is even then  $n$  is even.

Let  $p$  :  $3n + 2$  is even

$q$  :  $n$  is even

Let  $\sim q$  is true i.e.,  $n$  is odd

$$\Rightarrow n = 2k + 1, \text{ where } k \in Z$$

$$\text{Now, } 3n + 2 = 3(2k + 1) + 2$$

$$= 2(3k + 2) + 1$$

$$= 2m + 1 \text{ where } m = 3k + 2 \in Z$$

$$\therefore (3n + 2) \text{ is odd } \Rightarrow \sim p \text{ is true}$$

Hence  $\sim q \rightarrow \sim p$  is true.

$\therefore$  By method of contraposition  $p \rightarrow q$  is true.

- ii. **Proof by exhaustive cases** : Some proofs proceed by exhausting all the possibilities. We will examine a relatively small number of examples to prove the theorem. Such proofs are called exhaustive proofs.

**Example** : Prove that  $n^2 + 1 \geq 2^n$  where  $n$  is a positive integer and  $1 \leq n \leq 4$ .

We will verify the given inequality  $n^2 + 1 \geq 2^n$  for  $n = 1, 2, 3, 4$ .

For  $n = 1$ ;  $1 + 1 = 2$  which is true

For  $n = 2$ ;  $4 + 1 > 2^2$  which is true

For  $n = 3$ ;  $9 + 1 > 2^3$  which is true

For  $n = 4$ ;  $16 + 1 > 2^4$  which is true

In each of these four cases  $n^2 + 1 \geq 2^n$  holds true. Therefore, by method of exhaustive cases  $n^2 + 1 \geq 2^n$ , where  $n$  is the positive integer and  $1 \leq n \leq 4$  is true.

iii. **Proof by Cases :** In this method, we will cover up all the possible cases that we come across while proving the theorem.

**Example :** Prove that if  $x$  and  $y$  are real numbers, then  $\max(x, y) + \min(x, y) = x + y$ .

**Case I :** If  $x \leq y$ , then  $\max(x, y) + \min(x, y) = y + x = x + y$ .

**Case II :** if  $x \geq y$ , then  $\max(x, y) + \min(x, y) = x + y$ .

### VERY IMPORTANT QUESTIONS

*Following questions are very important. These questions may be asked in your SESSIONALS as well as UNIVERSITY EXAMINATION.*

**Q. 1.** Prove for any two sets  $A$  and  $B$  that,  $(A \cup B)' = A' \cap B'$ .

**ANS.** Refer Q. 1.8.

**Q. 2.** Show that  $R = \{(a, b) \mid a \equiv b \pmod{m}\}$  is an equivalence relation on  $Z$ . Show that if  $x_1 \equiv y_1$  and  $x_2 \equiv y_2$  then  $(x_1 + x_2) \equiv (y_1 + y_2)$ .

**ANS.** Refer Q. 1.18.

**Q. 3.** Let  $R$  be binary relation on the set of all strings of 0's and 1's such that  $R = \{(a, b) \mid a \text{ and } b \text{ are strings that have the same number of 0's}\}$ . Is  $R$  is an equivalence relation and a partial ordering relation ?

**ANS.** Refer Q. 1.19.

**Q. 4.** Let  $A = \{1, 2, 3, \dots, 13\}$ . Consider the equivalence relation on  $A \times A$  defined by  $(a, b) R (c, d)$  if  $a + d = b + c$ . Find equivalence classes of  $(5, 8)$ .

**ANS.** Refer Q. 1.20.

**Q. 5.** The following relation on  $A = \{1, 2, 3, 4\}$ . Determine whether the following :

a.  $R = \{(1, 3), (3, 1), (1, 1), (1, 2), (3, 3), (4, 4)\}$

b.  $R = A \times A$

Is an equivalence relation or not ?

**ANS.** Refer Q. 1.21.

**Q. 6.** Let  $n$  be a positive integer and  $S$  a set of strings. Suppose that  $R_n$  is the relation on  $S$  such that  $s R_n t$  if and only if  $s = t$ , or both  $s$  and  $t$  have at least  $n$  characters and first  $n$  characters of  $s$  and  $t$  are the same. That is, a string of fewer than  $n$  characters is related only to itself; a string  $s$  with at

least  $n$  characters is related to a string  $t$  if and only if  $t$  has at least  $n$  characters and  $t$  begins with the  $n$  characters at the start of  $s$ .

ANS Refer Q. 1.23.

Q. 7. Let  $X = \{1, 2, 3, \dots, 7\}$  and  $R = \{(x, y) \mid (x - y) \text{ is divisible by } 3\}$ . Is  $R$  equivalence relation. Draw the digraph of  $R$ .

ANS Refer Q. 1.24.

Q. 8. Determine whether each of these functions is a bijective from  $R$  to  $R$ .

a.  $f(x) = x^2 + 1$

b.  $f(x) = x^2$

c.  $f(x) = (x^2 + 1)/(x^2 + 2)$

ANS Refer Q. 1.27.

Q. 9. If  $f: A \rightarrow B$ ,  $g: B \rightarrow C$  are invertible functions, then show that  $g \circ f: A \rightarrow C$  is invertible and  $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$ .

ANS Refer Q. 1.28.

Q. 10. Prove that  $n^3 + 2n$  is divisible by 3 using principle of mathematical induction, where  $n$  is natural number.

ANS Refer Q. 1.32.

Q. 11. Prove by induction :  $\frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \dots + \frac{1}{n(n+1)} = \frac{n}{(n+1)}$ .

ANS Refer Q. 1.33.

Q. 12. Prove by the principle of mathematical induction, that the sum of finite number of terms of a geometric progression,

$$a + ar + ar^2 + \dots + ar^{n-1} = a(r^n - 1)/(r - 1) \text{ if } r \neq 1.$$

ANS Refer Q. 1.34.

Q. 13. Prove that if  $n$  is a positive integer, then 133 divides  $11^{n+1} + 12^{2n-1}$ .

ANS Refer Q. 1.35.

Q. 14. Prove by mathematical induction,  $n^4 - 4n^2$  is divisible by 3 for all  $n \geq 2$ .

ANS Refer Q. 1.36.

Q. 15. Prove by mathematical induction

$$3 + 33 + 333 + \dots + 3333 = (10^{n+1} - 9n - 10)/27$$

ANS Refer Q. 1.37.



# 2

UNIT

## Algebraic Structures

### CONTENTS

<b>Part-1</b> :	Algebraic Structures : .....	<b>2-2C to 2-10C</b>
	Definition, Groups, Subgroups and Order	
<b>Part-2</b> :	Cyclic Group, Cosets, .....	<b>2-10C to 2-16C</b>
	Lagrange's Theorem	
<b>Part-3</b> :	Normal Subgroups, .....	<b>2-17C to 2-18C</b>
	Permutation and Symmetric of Groups	
<b>Part-4</b> :	Group Homomorphisms, .....	<b>2-19C to 2-22C</b>
	Definition and Elementary Properties of Rings and Fields.	

## PART-1

Algebraic Structures : Definition, Groups, Subgroups and Order.

## Questions-Answers

Long Answer Type and Medium Answer Type Questions

**Que 2.1.** What is algebraic structure ? List properties of algebraic system.

**Answer**

**Algebraic structure :** An algebraic structure is a non-empty set  $G$  equipped with one or more binary operations. Suppose  $*$  is a binary operation on  $G$ . Then  $(G, *)$  is an algebraic structure.

**Properties of algebraic system :** Let  $(S, *, +)$  be an algebraic structure where  $*$  and  $+$  binary operation on  $S$  :

1. **Closure property :**  $(a * b) \in S \quad \forall a, b, \in S$
2. **Associative property :**  $(a * b) * c = a * (b * c) \quad \forall a, b, c \in S$
3. **Commutative property :**  $(a * b) = (b * a) \quad \forall a, b \in S$
4. **Identity element :**  $\exists e \in S$  such that  $a * e = a$  (right identity)  $\forall a \in S$ ,  $e$  is called identity element of  $S$  with respect to operation  $*$ .
5. **Inverse element :** For every  $a \in S$ ,  $\exists a^{-1} \in S$  such that  

$$a * a^{-1} = e = a^{-1} * a$$
 here  $a^{-1}$  is called inverse of 'a' under operation  $*$ .
6. **Cancellation property :**  
 $a * b = a * c \Rightarrow b = c$  and  $b * a = c * a \Rightarrow b = c \quad \forall a, b, c \in S$  and  $a \neq 0$
7. **Distributive property :**  $\forall a, b, c \in S$   
 $a * (b + c) = (a * b) + (a * c)$  (right distributive)  
 $(b + c) * a = (b * a) + (c * a)$  (left distributive)
8. **Idempotent property :** An element  $a \in S$  is called idempotent element with respect to operation  $*$  if  $a * a = a$ .

**Que 2.2.** Write short notes on :

- |                                |                    |
|--------------------------------|--------------------|
| i. Group                       | ii. Abelian group  |
| iii. Finite and infinite group | iv. Order of group |
| v. Groupoid                    |                    |

**Answer**

- i. **Group :** Let  $(G, *)$  be an algebraic structure where  $*$  is binary operation then  $(G, *)$  is called a group if following properties are satisfied :

1.  $a * b \in G \forall a, b \in G$  [closure property]
2.  $a * (b * c) = (a * b) * c \quad \forall a, b, c \in G$  [associative property]
3. There exist an element  $e \in G$  such that for any  $a \in G$   
 $a * e = e * a = e$  [existence of identity]
4. For every  $a \in G, \exists$  element  $a^{-1} \in G$   
such that  $a * a^{-1} = a^{-1} * a = e$

**For example :**  $(Z, +), (R, +),$  and  $(Q, +)$  are all groups.

ii. **Abelian group :** A group  $(G, *)$  is called abelian group or commutative group if binary operation  $*$  is commutative i.e.,  $a * b = b * a \quad \forall a, b \in G$

**For example :**  $(Z, +)$  is an abelian group.

iii. **Finite group :** A group  $(G, *)$  is called a finite group if number of elements in  $G$  are finite. For example  $G = \{0, 1, 2, 3, 4, 5\}$  under  $\otimes_6$  is a finite group.

**Infinite group :** A group  $(G, *)$  is called infinite group if number of element in  $G$  are infinite.

For example,  $(Z, +)$  is infinite group.

iv. **Order of group :** Order of group  $G$  is the number of elements in group  $G$ . It is denoted by  $o(G)$  or  $|G|$ . A group of order 1 has only the identity element.

v. **Groupoid :** Let  $(S, *)$  be an algebraic structure in which  $S$  is a non-empty set and  $*$  is a binary operation on  $S$ . Thus,  $S$  is closed with the operation  $*$ . Such a structure consisting of a non-empty set  $S$  and a binary operation defined in  $S$  is called a groupoid.

**Que 2.3.** Describe subgroup with example.

**Answer**

If  $(G, *)$  is a group and  $H \subseteq G$ . The  $(H, *)$  is said to subgroup of  $G$  if  $(H, *)$  is also a group by itself. i.e.,

(1)  $a * b \in H \quad \forall a, b \in H$  (Closure property)

(2)  $\exists e \in H$  such that  $a * e = a = e * a \quad \forall a \in H$

Where  $e$  is called identity of  $G$ .

(3)  $\exists a^{-1} \in H$  such that  $a * a^{-1} = e = a^{-1} * a \quad \forall a \in H$

**For example :** The set  $Q^+$  of all non-zero +ve rational number is subgroup of  $Q - \{0\}$ .

**Que 2.4.** Show that the set  $G = \{x + y\sqrt{2} \mid x, y \in Q\}$  is a group with respect to addition.

**Answer**

i. **Closure :**

$$\text{Let } X = x_1 + \sqrt{2} y_1$$

$$Y = x_2 + \sqrt{2} y_2$$

where  $x_1, x_2, y_1, y_2 \in Q$  and  $X, Y \in G$

$$\begin{aligned} \text{Then } X + Y &= (x_1 + \sqrt{2}y_1) + (x_2 + \sqrt{2}y_2) \\ &= (x_1 + x_2) + (y_1 + y_2)\sqrt{2} \\ &= X_1 + \sqrt{2}Y_1 \in G \text{ where } X_1, Y_1 \in \mathbb{Q} \end{aligned}$$

Therefore,  $G$  is closed under addition [ $\therefore$  Sum of two rational numbers is rational].

ii. **Associativity :**

Let  $X, Y$  and  $Z \in G$

$$\therefore \quad X = x_1 + \sqrt{2}y_1 \quad Y = x_2 + \sqrt{2}y_2 \quad Z = x_3 + \sqrt{2}y_3$$

where  $x_1, x_2, x_3, y_1, y_2, y_3 \in \mathbb{Q}$

$$\begin{aligned} \text{Consider } (X + Y) + Z &= (x_1 + \sqrt{2}y_1 + x_2 + \sqrt{2}y_2) + (x_3 + \sqrt{2}y_3) \\ &= ((x_1 + x_2) + (y_1 + y_2)\sqrt{2}) + (x_3 + \sqrt{2}y_3) \\ &= (x_1 + x_2 + x_3) + (y_1 + y_2 + y_3)\sqrt{2} \quad \dots(2.4.1) \end{aligned}$$

$$\begin{aligned} \text{Also } X + (Y + Z) &= (x_1 + \sqrt{2}y_1) + ((x_2 + \sqrt{2}y_2) + (x_3 + \sqrt{2}y_3)) \\ &= (x_1 + \sqrt{2}y_1) + ((x_2 + x_3) + (y_2 + y_3)\sqrt{2}) \\ &= (x_1 + x_2 + x_3) + (y_1 + y_2 + y_3)\sqrt{2} \quad \dots(2.4.2) \end{aligned}$$

From eq. (2.4.1) and (2.4.2)

$$(X + Y) + Z = X + (Y + Z)$$

Therefore,  $G$  is associative under addition.

iii. **Identity element :**

Let  $e \in G$  be identity elements of  $G$  under addition then

$$(x + y\sqrt{2}) + (e_1 + e_2\sqrt{2}) = x + y\sqrt{2}$$

where  $e = e_1 + e_2\sqrt{2}$  and  $e_1, e_2, x, y \in \mathbb{Q}$

$$\begin{aligned} e_1 + e_2\sqrt{2} &= 0 + 0\sqrt{2} \\ e_1 &= 0 \text{ and } e_2 = 0 \end{aligned}$$

Therefore,  $0 \in G$  is identity element.

iv. **Inverse element :**

$$-x - y\sqrt{2} \in G \text{ is inverse of } x + y\sqrt{2} \in G.$$

Therefore, inverse exist for every element  $x + y\sqrt{2} \in G$  such that,  $y \in \mathbb{Q}$ .

Hence,  $G$  is a group under addition.

**Que 2.5.** Let  $H$  be a subgroup of a finite group  $G$ . Prove that order

of  $H$  is a divisor of order of  $G$ .

AKTU 2018-19, Marks 07

**Answer**

- Let  $H$  be any sub-group of order  $m$  of a finite group  $G$  of order  $n$ . Let us consider the left coset decomposition of  $G$  relative to  $H$ .
- We will show that each coset  $aH$  consists of  $m$  different elements.  
Let  $H = \{h_1, h_2, \dots, h_m\}$

3. Then  $ah_1, ah_2, \dots, ah_m$ , are the members of  $aH$ , all distinct.  
For, we have

$$ah_i = ah_j \Rightarrow h_i = h_j$$

by cancellation law in  $G$ .

4. Since  $G$  is a finite group, the number of distinct left cosets will also be finite, say  $k$ . Hence the total number of elements of all cosets is  $km$  which is equal to the total number of elements of  $G$ .

Hence

$$n = mk$$

This show that  $m$ , the order of  $H$ , is a divisor of  $n$ , the order of the group  $G$ .

We also find that the index  $k$  is also a divisor of the order of the group.

**Que 2.6.** Define identity and zero elements of a set under a binary operation  $*$ . What do you mean by an inverse element ?

**Answer**

**Identity element :** An element  $e$  in a set  $S$  is called an identity element with respect to the binary operation  $*$  if, for any element  $a$  in  $S$

$$a * e = e * a = a$$

If  $a * e = a$ , then  $e$  is called the right identity element for the operation  $*$  and if  $e * a = a$ , then  $e$  is called the left identity element for the operation  $*$ .

**Zero element :** Let  $R$  be an abelian group with respect to addition. The element  $0 \in R$  will be the additive identity. It is called the zero element of  $R$ .

**Inverse element :** Consider a set  $S$  having the identity element  $e$  with respects to the binary operation  $*$ . If corresponding to each element  $a \in S$  there exists an element  $b \in S$  such that

$$a * b = b * a = e$$

Then  $b$  is said to be the inverse of  $a$  and is usually denoted by  $a^{-1}$ . We say  $a$  is invertible.

**Que 2.7.** Prove that  $(Z_6, (+_6))$  is an abelian group of order 6, where

$$Z_6 = \{0, 1, 2, 3, 4, 5\}.$$

**AKTU 2014-15, Marks 05**

**Answer**

The composition table is :

$+_6$	0	1	2	3	4	5
0	0	1	2	3	4	5
1	1	2	3	4	5	0
2	2	3	4	5	0	1
3	3	4	5	0	1	2
4	4	5	0	1	2	3
5	5	0	1	2	3	4

Since

$$2 +_6 1 = 3$$

$$4 +_6 5 = 3$$

From the table we get the following observations :

**Closure :** Since all the entries in the table belong to the given set  $Z_6$ . Therefore,  $Z_6$  is closed with respect to addition modulo 6.

**Associativity :** The composition '+<sub>6</sub>' is associative. If  $a, b, c$  are any three elements of  $Z_6$ ,

$$\begin{aligned} a +_6 (b +_6 c) &= a +_6 (b + c) \quad [\because b +_6 c = b + c \pmod{6}] \\ &= \text{least non-negative remainder when } a + (b + c) \text{ is divided by 6.} \\ &= \text{least non-negative remainder when } (a + b) + c \text{ is divided by 6.} \\ &= (a + b) +_6 c = (a +_6 b) +_6 c. \end{aligned}$$

**Identity :** We have  $0 \in Z_6$ . If  $a$  is any element of  $Z_6$ , then from the composition table we see that

$$0 +_6 a = a = a +_6 0$$

Therefore, 0 is the identity element.

**Inverse :** From the table we see that the inverse of 0, 1, 2, 3, 4, 5 are 0, 5, 4, 3, 2, 1 respectively. For example  $4 +_6 2 = 0 = 2 +_6 4$  implies 4 is the inverse of 2.

**Commutative :** The composition is commutative as the elements are symmetrically arranged about the main diagonal. The number of elements in the set  $Z_6$  is 6.

$\therefore (Z_6, +_6)$  is a finite abelian group of order 6.

**Que 2.8.** Let  $G = \{1, -1, i, -i\}$  with the binary operation

multiplication be an algebraic structure, where  $i = \sqrt{-1}$ . Determine whether  $G$  is an abelian or not.

AKTU 2018-19, Marks 07

**Answer**

The composition table of  $G$  is

*	1	-1	i	-i
1	1	-1	i	-i
-1	-1	1	-i	i
i	i	-i	-1	1
-i	-i	i	-1	1

- Closure property :** Since all the entries of the composition table are the elements of the given set, the set  $G$  is closed under multiplication.
- Associativity :** The elements of  $G$  are complex numbers, and we know that multiplication of complex numbers is associative.
- Identity :** Here, 1 is the identity element.
- Inverse :** From the composition table, we see that the inverse elements of 1, -1, i, -i are 1, -1, -i, i respectively.

5. **Commutativity** : The corresponding rows and columns of the table are identical. Therefore the binary operation is commutative. Hence,  $(G, *)$  is an abelian group.

**Que 2.9.** Write the properties of group. Show that the set  $\{1, 2, 3, 4, 5\}$  is not group under addition and multiplication modulo 6.

AKTU 2017-18, Marks 07

**Answer**

Properties of group : Refer Q. 2.2(i). Page 2-2C, Unit-2.

Numerical :

**Addition modulo 6 ( $+_6$ )** : Composition table of  $S = \{1, 2, 3, 4, 5\}$  under operation  $+_6$  is given as :

$+_6$	1	2	3	4	5
1	2	3	4	5	0
2	3	4	5	0	1
3	4	5	0	1	2
4	5	0	1	2	3
5	0	1	2	3	4

Since,  $1 +_6 5 = 0$  but  $0 \notin S$  i.e.,  $S$  is not closed under addition modulo 6. So,  $S$  is not a group.

**Multiplication modulo 6 ( $*_6$ )** :

Composition table of  $S = \{1, 2, 3, 4, 5\}$  under operation  $*_6$  is given as

$*_6$	1	2	3	4	5
1	1	2	3	4	5
2	2	4	0	2	4
3	3	0	3	0	3
4	4	2	0	4	2
5	5	4	3	2	1

Since,  $2 *_6 3 = 0$  but  $0 \notin S$  i.e.,  $S$  is not closed under multiplication modulo 6. So,  $S$  is not a group.

**Que 2.10.** Let  $G = \{a, a^2, a^3, a^4, a^5, a^6 = e\}$ . Find the order of every element.

**Answer**

$$o(a) : a^6 = e \rightarrow o(a) = 6$$

$$o(a^2) : (a^2)^3 = a^6 = e \rightarrow o(a^2) = 3$$

$$o(a^3) : (a^3)^2 = a^6 = e \rightarrow o(a^3) = 2$$

$$o(a^4) : (a^4)^3 = a^{12} = (a^6)^2 = e^2 = e \rightarrow o(a^4) = 3$$

$$o(a^5) : (a^5)^6 = a^{30} = (a^6)^5 = e^5 = e \rightarrow o(a^5) = 6$$

$$o(a^6) : (a^6)^1 = a^6 = e \rightarrow o(a^6) = 1$$

**Que 2.11.** Let  $G$  be a group and let  $a, b \in G$  be any elements.

Then

- i.  $(a^{-1})^{-1} = a$
- ii.  $(a * b)^{-1} = b^{-1} * a^{-1}$ .

AKTU 2014-15, Marks 05

Answer

- i. Let  $e$  be the identity element for  $*$  in  $G$ .  
Then we have  $a * a^{-1} = e$ , where  $a^{-1} \in G$ .  
Also  $(a^{-1})^{-1} * a^{-1} = e$   
Therefore,  $(a^{-1})^{-1} * a^{-1} = a * a^{-1}$ .  
Thus, by right cancellation law, we have  $(a^{-1})^{-1} = a$ .
- ii. Let  $a$  and  $b \in G$  and  $G$  is a group for  $*$ , then  $a * b \in G$  (closure)  
Therefore,  $(a * b)^{-1} * (a * b) = e$ . ....(2.11.1)  
Let  $a^{-1}$  and  $b^{-1}$  be the inverses of  $a$  and  $b$  respectively, then  $a^{-1}, b^{-1} \in G$ .  
Therefore,  $(b^{-1} * a^{-1}) * (a * b) = b^{-1} * (a^{-1} * a) * b$  (associativity)  
 $= b^{-1} * e * b = b^{-1} * b = e$  ....(2.11.2)  
From (2.11.1) and (2.11.2) we have,  
 $(a * b)^{-1} * (a * b) = (b^{-1} * a^{-1}) * (a * b)$   
 $(a * b)^{-1} = b^{-1} * a^{-1}$  (by right cancellation law)

**Que 2.12.** Prove that the intersection of two subgroups of a group

is also subgroup.

AKTU 2014-15, Marks 05

Answer

Let  $H_1$  and  $H_2$  be any two subgroups of  $G$ . Since at least the identity element  $e$  is common to both  $H_1$  and  $H_2$ .

$$\therefore H_1 \cap H_2 \neq \phi$$

In order to prove that  $H_1 \cap H_2$  is a subgroup, it is sufficient to prove that

$$a \in H_1 \cap H_2, b \in H_1 \cap H_2 \Rightarrow ab^{-1} \in H_1 \cap H_2$$

Now  $a \in H_1 \cap H_2 \Rightarrow a \in H_1$  and  $a \in H_2$

$$b \in H_1 \cap H_2 \Rightarrow b \in H_1 \text{ and } b \in H_2$$

But  $H_1, H_2$  are subgroups. Therefore,

$$a \in H_1, b \in H_1 \Rightarrow ab^{-1} \in H_1$$

$$a \in H_2, b \in H_2 \Rightarrow ab^{-1} \in H_2$$

Finally,  $ab^{-1} \in H_1, ab^{-1} \in H_2 \Rightarrow ab^{-1} \in H_1 \cap H_2$

Thus, we have shown that

$$a \in H_1 \cap H_2, b \in H_1 \cap H_2 \Rightarrow ab^{-1} \in H_1 \cap H_2.$$

Hence,  $H_1 \cap H_2$  is a subgroup of  $G$ .

**Que 2.13.** Let  $G$  be the set of all non-zero real number and let  $a * b = ab/2$ . Show that  $(G, *)$  be an abelian group.

AKTU 2015-16, Marks 10

**Answer**

i. **Closure property :** Let  $a, b \in G$ .

$$a * b = \frac{ab}{2} \in G \text{ as } ab \neq 0$$

$\Rightarrow *$  is closure in  $G$ .

ii. **Associativity :** Let  $a, b, c \in G$

$$\text{Consider } a * (b * c) = a * \left(\frac{bc}{2}\right) = \frac{a(bc)}{4} = \frac{abc}{4}$$

$$(a * b) * c = \left(\frac{ab}{2}\right) * c = \frac{(ab)c}{4} = \frac{abc}{4}$$

$\Rightarrow *$  is associative in  $G$ .

iii. **Existence of the identity :** Let  $a \in G$  and  $\exists e$  such that

$$a * e = \frac{ae}{2} = a$$

$$\Rightarrow ae = 2a$$

$$\Rightarrow e = 2$$

$\therefore 2$  is the identity element in  $G$ .

iv. **Existence of the inverse :** Let  $a \in G$  and  $b \in G$  such that  $a * b = e = 2$

$$\Rightarrow \frac{ab}{2} = 2$$

$$\Rightarrow ab = 4$$

$$\Rightarrow b = \frac{4}{a}$$

$\therefore$  The inverse of  $a$  is  $\frac{4}{a}, \forall a \in G$ .

v. **Commutative :** Let  $a, b \in G$

$$a * b = \frac{ab}{2}$$

$$\text{and } b * a = \frac{ba}{2} = \frac{ab}{2}$$

$\Rightarrow *$  is commutative.

Thus,  $(G, *)$  is an abelian group.

**Que 2.14.** Prove that inverse of each element in a group is unique.

AKTU 2015-16, Marks 10

**Answer**

Let (if possible)  $b$  and  $c$  be two inverses of element  $a \in G$ .  
Then by definition of group :

$$b * a = a * b = e$$

and

$$a * c = c * a = e$$

where  $e$  is the identity element of  $G$

Now

$$\begin{aligned} b &= e * b = (c * a) * b \\ &= c * (a * b) \\ &= c * e \\ &= c \\ b &= c \end{aligned}$$

Therefore, inverse of an element is unique in  $(G, *)$ .

**PART-2**

*Cyclic Group, Cosets, Lagrange's Theorem.*

**Questions-Answers****Long Answer Type and Medium Answer Type Questions**

**Que 2.15.** Define cyclic group with suitable example.

**Answer**

**Cyclic group :** A group  $G$  is called a cyclic group if  $\exists$  at least one element  $a$  in  $G$  such that every element  $x \in G$  is of the form  $a^n$ , where  $n$  is some integer. The element  $a \in G$  is called the generator of  $G$ .

**For example :**

Show that the multiplicative group  $G = \{1, -1, i, -i\}$  is cyclic. Also find its generators.

We have,

$$i^1 = i, i^2 = -1, i^3 = -i, i^4 = 1.$$

and

$$(-i)^1 = -i, (-i)^2 = -1, (-i)^3 = i, (-i)^4 = 1$$

Thus, every element in  $G$  be expressed as  $i^n$  or  $(-i)^n$

$\therefore G$  is cyclic group and its generators are  $i$  and  $-i$ .

**Que 2.16.** Prove that every group of prime order is cyclic.

AKTU 2018-19, Marks 07

**Answer**

1. Let  $G$  be a group whose order is a prime  $p$ .
2. Since  $P > 1$ , there is an element  $a \in G$  such that  $a \neq e$ .

3. The group  $\langle a \rangle$  generated by 'a' is a subgroup of  $G$ .
4. By Lagrange's theorem, the order of  $\langle a \rangle$  divides  $|G|$ .
5. But the only divisors of  $|G| = p$  are 1 and  $p$ . Since  $a \neq e$  we have  $|\langle a \rangle| > 1$ , so  $|\langle a \rangle| = p$ .
6. Hence,  $\langle a \rangle = G$  and  $G$  is cyclic.

**Que 2.17.** Show that every group of order 3 is cyclic.

**AKTU 2014-15, Marks 05**

**Answer**

1. Suppose  $G$  is a finite group whose order is a prime number  $p$ , then to prove that  $G$  is a cyclic group.
2. An integer  $p$  is said to be a prime number if  $p \neq 0$ ,  $p \neq \pm 1$ , and if the only divisors of  $p$  are  $\pm 1$ ,  $\pm p$ .
3. Some  $G$  is a group of prime order, therefore  $G$  must contain at least 2 element. Note that 2 is the least positive prime integer.
4. Therefore, there must exist an element  $a \in G$  such that  $a \neq$  the identity element  $e$ .
5. Since  $a$  is not the identity element, therefore  $o(a)$  is definitely  $\geq 2$ . Let  $o(a) = m$ . If  $H$  is the cyclic subgroup of  $G$  generated by  $a$  then  $o(H) = o(a) = m$ .
6. By Lagrange's theorem  $m$  must be a divisor of  $p$ . But  $p$  is prime and  $m \geq 2$ . Hence,  $m = p$ .
7.  $\therefore H = G$ . Since  $H$  is cyclic therefore  $G$  is cyclic and  $a$  is a generator of  $G$ .

**Que 2.18.** Show that group  $(G, +_5)$  is a cyclic group where  $G = \{0, 1, 2, 3, 4\}$ . What are its generators ?

**Answer**

Composition table of  $G$  under operation  $+_5$  is shown in Table 2.18.1.

**Table 2.18.1.**

$+_5$	0	1	2	3	4
0	0	1	2	3	4
1	1	2	3	4	0
2	2	3	4	0	1
3	3	4	0	1	2
4	4	0	1	2	3

0 is the identity element of  $G$  under  $+_5$ .

$$1^1 = 1 \equiv 1 \pmod{5}$$

$$1^2 = 1 + 1 \equiv 2 \pmod{5}$$

$$1^3 = 1 + 1 + 1 \equiv 3 \pmod{5}$$

$$1^4 = 1 + 1 + 1 + 1 \equiv 4 \pmod{5}$$

$$1^5 = 1 + 1 + 1 + 1 + 1 \equiv 0 \pmod{5}$$

Downloaded from

where  $1^n$  means 1 is added  $n$  times

Therefore, 1 is generator of  $G$ .

Similarly, 2, 3, 4 are also generators of  $G$ .

Therefore,  $G$  is a cyclic group with generator 1, 2, 3, 4.

**Que 2.19.** Show that  $G = [(1, 2, 4, 5, 7, 8), X_9]$  is cyclic. How many

generators are there? What are they? **AKTU 2015-16, Marks 7.5**

**Answer**

Composition table for  $X_9$  is

$X_9$	1	2	4	5	7	8
1	2	3	4	5	7	8
2	2	4	8	1	5	7
4	4	8	7	2	1	5
5	5	1	2	7	8	4
7	7	5	1	8	4	2
8	8	7	5	4	2	1

1 is identity element of group  $G$

$$2^1 = 2 = 2 \pmod{9}$$

$$2^2 = 4 = 4 \pmod{9}$$

$$2^3 = 8 = 8 \pmod{9}$$

$$2^4 = 16 = 7 \pmod{9}$$

$$2^5 = 32 = 5 \pmod{9}$$

$$2^6 = 64 = 1 \pmod{9}$$

Therefore, 2 is generator of  $G$ . Hence  $G$  is cyclic.

Similarly, 5 is also generator of  $G$ .

Hence there are two generators 2 and 5.

**Que 2.20.** Define cosets. Write and prove properties of cosets.

**Answer**

Let  $H$  be a subgroup of group  $G$  and let  $a \in G$  then the set  $Ha = \{ha : h \in H\}$  is called right coset generated by  $H$  and  $a$ .

Also the set  $aH = \{ah : h \in H\}$  is called left coset generated by  $a$  and  $H$ .

**Properties of cosets:** Let  $H$  be a subgroup of  $G$  and let  $a$  and  $b$  belong to  $G$ . Then

1.  $a \in aH$

**Proof:**  $a = ae \in aH$

Since  $e$  is identity element of  $G$ .

2.  $aH = H$  iff  $a \in H$ .

**Proof:** Let  $aH = H$ .

Then  $a = ae \in aH = H$  ( $e$  is identity in  $G$  and so is in  $H$ )

$\Rightarrow a \in H$

3.  $aH = bH$  or  $aH \cap bH = \emptyset$

**Proof :** Let  $aH = bH$  or  $aH \cap bH = \phi$   
and to prove that  $aH = bH$ .

Let  $aH \cap bH$

Then there exists  $h_1, h_2 \in H$  such that

$$x = ah_1 \text{ and } x = bh_2$$

$$a = xh_1^{-1} = bh_2 h_1^{-1}$$

Since  $H$  is a subgroup, we have  $h_2 h_1^{-1} \in H$

$$\text{let } h_2 h_1^{-1} = h \in H$$

Now,  $aH = bh_2 h_1^{-1} H = (bh) H = b(hH) = bh (\because hH = H \text{ by property 2})$

$$\therefore aH = bH \text{ if } aH \cap bH \neq \phi$$

Thus, either  $aH \cap bH = \phi$  or  $aH = bH$ .

4.  $aH = bH$  iff  $a^{-1}b \in H$ .

**Proof :** Let  $aH = bH$ .

$$a^{-1}aH = a^{-1}bH$$

$$eH = a^{-1}bH$$

$$H = (a^{-1}b)H$$

( $e$  is identity in  $G$ )

Therefore by property (2);  $a^{-1}b \in H$ .

Conversely, now if  $a^{-1}b \in H$ .

Then consider  $bH = e(bH) = (a a^{-1})(bH)$

$$= a(a^{-1}b)H$$

$$= aH$$

Thus  $aH = bH$  iff  $a^{-1}b \in H$ .

5.  $aH$  is a subgroup of  $G$  iff  $a \in H$ .

**Proof :** Let  $aH$  is a subgroup of  $G$  then it contains the identity  $e$  of  $G$ .

Thus,  $aH \cap eH \neq \phi$

then by property (3);  $aH = eH = H$

$$aH = H \Rightarrow a \in H$$

Conversely, if  $a \in H$  then by property (2);  $aH = H$ .

**Que 221.** State and prove Lagrange's theorem for group. Is the

converse true ?

**AKTU 2016-17, Marks 10**

**Answer**

**Lagrange's theorem :**

**Statement :** The order of each subgroup of a finite group is a divisor of the order of the group.

**Proof :** Let  $G$  be a group of finite order  $n$ . Let  $H$  be a subgroup of  $G$  and let  $O(H) = m$ . Suppose  $h_1, h_2, \dots, h_m$  are the  $m$  members of  $H$ .

Let  $a \in G$ , then  $Ha$  is the right coset of  $H$  in  $G$  and we have

$$Ha = \{h_1 a, h_2 a, \dots, h_m a\}$$

$Ha$  has  $m$  distinct members, since  $h_i a = h_j a \Rightarrow h_i = h_j$

Therefore, each right coset of  $H$  in  $G$  has  $m$  distinct members. Any two distinct right cosets of  $H$  in  $G$  are disjoint i.e., they have no element in

common. Since  $G$  is a finite group, the number of distinct right cosets of  $H$  in  $G$  will be finite, say, equal to  $k$ . The union of these  $k$  distinct right cosets of  $H$  in  $G$  is equal to  $G$ .

Thus, if  $Ha_1, Ha_2, \dots, Ha_k$  are the  $k$  distinct right cosets of  $H$  in  $G$ . Then

$$G = Ha_1 \cup Ha_2 \cup Ha_3 \cup \dots \cup Ha_k$$

$\Rightarrow$  the number of elements in  $G =$  the number of elements in  $Ha_1 + \dots +$  the number of elements in  $Ha_2 + \dots +$  the number of elements in  $Ha_k$

$$\Rightarrow O(G) = km$$

$$\Rightarrow n = km$$

$$\Rightarrow k = \frac{n}{m}$$

$\Rightarrow m$  is a divisor of  $n$ .

$\Rightarrow O(H)$  is a divisor of  $O(G)$ .

**Proof of converse :** If  $G$  be a finite group of order  $n$  and  $n \in G$ , then

$$a^n = e$$

Let  $o(a) = m$  which implies  $a^m = e$ .

Now, the subset  $H$  of  $G$  consisting of all the integral power of  $a$  is a subgroup of  $G$  and the order of  $H$  is  $m$ .

Then, by the Lagrange's theorem,  $m$  is divisor of  $n$ .

Let  $n = mk$ , then

$$a^n = a^{mk} = (a^m)^k = e^k = e$$

$\therefore$  Yes, the converse is true.

**Que 2.22.** State and explain Lagrange's theorem.

**Answer**

**Lagrange's theorem :**

If  $G$  is a finite group and  $H$  is a subgroup of  $G$  then  $o(H)$  divides  $o(G)$ .

Moreover, the number of distinct left (right) cosets of  $H$  in  $G$  is  $o(G)/o(H)$ .

**Proof :** Let  $H$  be subgroup of order  $m$  of a finite group  $G$  of order  $n$ .

Let  $H = \{h_1, h_2, \dots, h_m\}$

Let  $a \in G$ . Then  $aH$  is a left coset of  $H$  in  $G$  and  $aH = \{ah_1, ah_2, \dots, ah_m\}$  has  $m$  distinct elements as  $ah_i = ah_j \Rightarrow h_i = h_j$  by cancellation law in  $G$ .

Thus, every left coset of  $H$  in  $G$  has  $m$  distinct elements.

Since  $G$  is a finite group, the number of distinct left cosets will also be finite.

Let it be  $k$ . Then the union of these  $k$ -left cosets of  $H$  in  $G$  is equal to  $G$ .

i.e., if  $a_1H, a_2H, \dots, a_kH$  are right cosets of  $H$  in  $G$  then

$$G = a_1H \cup a_2H \cup \dots \cup a_kH.$$

$$\therefore o(G) = o(a_1H) + o(a_2H) + \dots + o(a_kH)$$

(Since two distinct left cosets are mutually disjoint.)

$$\Rightarrow n = m + m + \dots + m \text{ (k times)}$$

$$\Rightarrow n = mk \Rightarrow k = \frac{n}{m}$$

$$\therefore k = \frac{o(G)}{o(H)}$$

Thus order of each subgroup of a finite group  $G$  is a divisor of the order of the group.

**Cor 1 :** If  $H$  has  $m$  different cosets in  $G$  then by Lagrange's theorem :

$$o(G) = m \cdot o(H)$$

$$\Rightarrow m = \frac{o(G)}{o(H)}$$

$$\therefore [G : H] = \frac{o(G)}{o(H)}$$

**Cor 2 :** If  $|G| = n$  and  $a \in G$  then  $a^n = e$

Let  $|a| = m \Rightarrow a^m = e$

Now, the subset  $H$  of  $G$  consisting of all integral powers of  $a$  is a subgroup of  $G$  and the order of  $H$  is  $m$ .

Then by Lagrange's theorem,  $m$  is divisor of  $n$ .

Let  $n = mk$ , then  $a^n = a^{mk} = (a^m)^k = e^k = e$

**Que 2.23.**

- Prove that every cyclic group is an abelian group.
- Obtain all distinct left cosets of  $\{(0), (3)\}$  in the group  $(Z_6, +_6)$  and find their union.
- Find the left cosets of  $\{[0], [3]\}$  in the group  $(Z_6, +_6)$ .

AKTU 2016-17, Marks 10

**Answer**

a. Let  $G$  be a cyclic group and let  $a$  be a generator of  $G$  so that  $G = \langle a \rangle = \{a^n : n \in Z\}$

If  $g_1$  and  $g_2$  are any two elements of  $G$ , there exist integers  $r$  and  $s$  such that  $g_1 = a^r$  and  $g_2 = a^s$ . Then

$$g_1 g_2 = a^r a^s = a^{r+s} = a^{s+r} = a^s \cdot a^r = g_2 g_1$$

So,  $G$  is abelian.

b.  $\therefore [0] + H = [3] + H, [1] + [4] + H$  and  $[2] + H = [5] + H$  are the three distinct left cosets of  $H$  in  $(Z_6, +_6)$ .

We would have the following left cosets :

$$\begin{aligned} g_1 H &= \{g_1 h, h \in H\} \\ g_2 H &= \{g_2 h, h \in H\} \\ g_n H &= \{g_n h, h \in H\} \end{aligned}$$

The union of all these sets will include all the  $g'$  s, since for each set

we have  $g_r \in g_k = \{g_k h, h \in H\}$  where  $e$  is the identity.

Then if we make the union of all these sets we will have at least all the elements of  $G$ . The other elements are merely  $g_h$  for some  $h$ . But since  $g_h \in G$  they would be repeated elements in the union. So, the union of all left cosets of  $H$  in  $G$  is  $G$ , i.e.,

$$Z_6 = \{[0], [1], [2], [3], [4], [5]\}$$

c. Let  
The le

**Que 2.24.**

$G = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$   
If  $H$  is a s  
 $H$  and its

**Answer**

Lagrange  
Numerica

Left coset o

**Que 2.25.**

group  $Z$   
 $H = \{\dots, -10, -5, 0, 5, 10, \dots\}$   
i. Find  
ii. What

**Answer**

Coset : Re  
Numerica  
i. We ha  
and  
Let 0

Now, i  
 $H, H +$   
disjoin  
Index o  
Theref

c. Let

 $Z_6 = \{[0], [1], [2], [3], [4], [5]\}$  be a group. $H = \{[0], [3]\}$  be a subgroup of  $(Z_6, +_6)$ .The left cosets of  $H$  are,

$$[0] + H = \{[0], [3]\}$$

$$[1] + H = \{[1], [4]\}$$

$$[2] + H = \{[2], [5]\}$$

$$[3] + H = \{[3], [0]\}$$

$$[4] + H = \{[4], [1]\}$$

$$[5] + H = \{[5], [2]\}$$

**Que 2.24.**

Write and prove the Lagrange's theorem. If a group  $G = \{\dots, -3, 2, -1, 0, 1, 2, 3, \dots\}$  having the addition as binary operation. If  $H$  is a subgroup of group  $G$  where  $x^2 \in H$  such that  $x \in G$ . What is  $H$  and its left coset w.r.t 1?

**AKTU 2014-15, Marks 05****Answer**

Lagrange's theorem : Refer Q. 2.22, Page 2-14C, Unit-2.

Numerical :

$$H = \{x^2 : x \in G\} = \{0, 1, 4, 9, 16, 25, \dots\}$$

Left coset of  $H$  will be  $1 + H = \{1, 2, 5, 10, 17, 26, \dots\}$ **Que 2.25.**

What do you mean by cosets of subgroup? Consider the group  $Z$  of integers under addition and the subgroup  $H = \{\dots, -10, -5, 0, 5, 10, \dots\}$  considering the multiple of 5.

i. Find the cosets of  $H$  in  $Z$ .ii. What is index of  $H$  in  $Z$ ?**Answer**

Coset : Refer Q. 2.20, Page 2-12C, Unit-2.

Numerical :

i. We have  $Z = \{-7, -6, -5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, \dots\}$ and  $H = \{\dots, -10, -5, 0, 5, 10, \dots\}$ Let  $0 \in Z$  and the right cosets are given as

$$H = H + 0 = \{\dots, -10, -5, 0, 5, 10, \dots\}$$

$$H + 1 = \{\dots, -9, -4, 1, 6, 11, \dots\}$$

$$H + 2 = \{\dots, -8, -3, 2, 7, 12, \dots\}$$

$$H + 3 = \{\dots, -7, -2, 3, 8, 13, \dots\}$$

$$H + 4 = \{\dots, -6, -1, 4, 9, 14, \dots\}$$

$$H + 5 = \{\dots, -10, -5, 0, 5, 10, \dots\} = H$$

Now, its repetition starts. Now, we see that the right cosets,  $H, H + 1, H + 2, H + 3, H + 4$  are all distinct and more over they are disjoint. Similarly the left cosets will be same as right cosets.

ii. Index of  $H$  in  $G$  is the number of distinct right/left cosets.

Therefore, index is 5.

## PART-3

Normal Subgroups, Permutation and Symmetric of Groups.

## Questions-Answers

Long Answer Type and Medium Answer Type Questions

**Que 2.26.** Write short notes on :

- Normal subgroup
- Permutation group

**Answer**

- Normal subgroup** : A subgroup  $H$  of  $G$  is said to be normal subgroup of  $G$  if  $Ha = aH \forall a \in G$  i.e. the right coset and left coset of  $H$  is generated by  $a$  are the same.
  - Clearly, every subgroup  $H$  of an abelian group  $G$  is a normal subgroup of  $G$ . For  $a \in G$  and  $h \in H$ ,  $ah = ha$ .
  - Since a cyclic group is abelian, every subgroup of a cyclic group is normal.
- Permutation group** : A set  $G$  of all permutation on a non-empty set  $A$  under the binary operation  $*$  is called permutation group. If  $A = \{1, 2, 3, \dots, n\}$ , the given permutation group formed by  $A$  is also called symmetric group of degree  $n$  denoted by  $S_n$ . Number of elements of  $S_n$  will be  $n!$ .

**Cyclic permutation** : Let  $A = \{x_1, x_2, \dots, x_n\}$ . Then let  $t_1, t_2, \dots, t_k$  be elements of set  $A$  and permutation  $P : A \rightarrow A$  defined by

$$P(t_1) = t_2$$

$$P(t_2) = t_3$$

.....

.....

$$P(t_{k-1}) = t_k$$

$$P(t_k) = t_1$$

is called a cyclic permutation of length  $k$ .

**For example** : Consider  $A = \{a, b, c, d, e\}$ . Then let  $P = \begin{pmatrix} a & b & c & d & e \\ c & b & d & a & e \end{pmatrix}$ .

Then  $P$  has a cycle of length 3 given by  $(a, c, d)$ .

**Que 2.27.** Define the subgroup of a group. Let  $(G, o)$  be a group. Let  $H = \{a \mid a \in G \text{ and } a \circ b = b \circ a \text{ for all } b \in G\}$ . Show that  $H$  is a normal subgroup.

**Answer**

**Subgroup :** If  $(G, *)$  is a group and  $H \subseteq G$ . Then  $(H, *)$  is said to subgroup of  $G$  if  $(H, *)$  is also a group by itself.

i.e.,

1.  $a * b \in H \forall a, b \in H$  (Closure property)
2.  $\exists e \in H$  such that  $a * e = a = e * a \quad \forall a \in H$   
where  $e$  is called identity of  $G$ .
3.  $\exists a^{-1} \in H$  such that  $a * a^{-1} = e = a^{-1} * a \quad \forall a \in H$

**Numerical :** Let  $(G, \circ)$  be a group. A non-empty subset  $H$  of a group  $G$  is said to be a subgroup of  $G$  if  $(H, \circ)$  itself is a group.

Given that,  $H = \{a \mid a \in G \text{ and } a \circ b = b \circ a; \forall b \in G\}$

Let  $a, b \in H \Rightarrow a \circ x = x \circ a$  and  $b \circ x = x \circ b, \forall x \in G$

$$\Rightarrow (b \circ x)^{-1} = (x \circ b)^{-1}$$

$$\Rightarrow x^{-1} \circ b^{-1} = b^{-1} \circ x^{-1}$$

$$\Rightarrow b^{-1} \in H.$$

$$\begin{aligned} \text{Now, } (a \circ b^{-1}) \circ x &= a \circ (b^{-1} \circ x) && [\because \circ \text{ is associative}] \\ &= a \circ (x \circ b^{-1}) && [\because \text{ use } b^{-1} \in H] \\ &= (a \circ x) \circ b^{-1} && [\because a \in H] \\ &= (x \circ a) \circ b^{-1} \\ &= x \circ (a \circ b^{-1}) \end{aligned}$$

$$\Rightarrow a \circ b^{-1} \in H$$

Therefore,  $H$  is a subgroup of group  $G$ .

Let  $h \in H$  and  $g \in G$  and any  $x$  in  $G$ .

Consider

$$\begin{aligned} (g \circ h \circ g^{-1}) \circ x &= (g \circ g^{-1} \circ h) \circ x && [\because h \in H] \\ &= (e \circ h) \circ x = h \circ x \\ &= x \circ h && [\because h \in H] \\ &= x \circ (h \circ g \circ g^{-1}) \\ &= x \circ (g \circ h \circ g^{-1}) && [\because h \in H] \end{aligned}$$

$$\Rightarrow g \circ h \circ g^{-1} \in H \text{ for any } g \in G$$

$\therefore H$  is a normal subgroup of  $G$ .

**Que 2.28.** If  $N$  and  $M$  are normal subgroup of  $G$  then  $N \cap M$  is a normal subgroup of  $G$ .

**Answer**

As  $N$  and  $M$  are subgroups of  $G$  then  $N \cap M$  is a subgroup of  $G$ . Let  $g \in G$  and  $a \in N \cap M$

$a \in N$  and  $a \in M$ .

Since  $N$  is normal subgroup of  $G$ ,  $gag^{-1} \in N$

Since  $M$  is normal subgroup of  $G$ ,  $gag^{-1} \in M$

$\therefore gag^{-1} \in N \cap M$  is a normal subgroup of  $G$ .

Hence  $N \cap M$  is a normal subgroup of  $G$ .

## PART-4

## Group Homomorphism, Definition and Elementary Properties of Ring and Fields.

## Questions-Answers

## Long Answer Type and Medium Answer Type Questions

**Que 2.29.** Discuss homomorphism and isomorphic group.

**Answer**

**Homomorphism :** Let  $(G_1, \cdot)$  and  $(G_2, *)$  be two groups then a mapping  $f: G_1 \rightarrow G_2$  is called a homomorphism if  $f(a \cdot b) = f(a) * f(b)$  for all  $a, b \in G_1$ . Thus  $f$  is homomorphism from  $G_1$  to  $G_2$  then  $f$  preserves the composition in  $G_1$  and  $G_2$  i.e., image of composition is equal to composition of images.

The group  $G_2$  is said to be homomorphic image of group  $G_1$  if there exist a homomorphism of  $G_1$  onto  $G_2$ .

**Isomorphism :** Let  $(G_1, \cdot)$  and  $(G_2, *)$  be two groups then a mapping  $f: G_1 \rightarrow G_2$  is an isomorphism if

- i.  $f$  is homomorphism.
- ii.  $f$  is one to one i.e.,  $f(x) = f(y) \Rightarrow x = y \quad \forall x, y \in G_1$ .
- iii.  $f$  is onto.

**Que 2.30.** Give the definitions of rings, integral domains and fields.

**Answer**

**Ring :** A ring is an algebraic system  $(R, +, \cdot)$  where  $R$  is a non-empty set and  $+$  and  $\cdot$  are two binary operations (which can be different from addition and multiplication) and if the following conditions are satisfied :

1.  $(R, +)$  is an abelian group.
2.  $(R, \cdot)$  is semigroup i.e.,  $(a \cdot b) \cdot c = a \cdot (b \cdot c) \quad \forall a, b, c \in R$ .
3. The operation  $\cdot$  is distributive over  $+$ .

i.e., for any  $a, b, c \in R$

$$a \cdot (b + c) = (a \cdot b) + (a \cdot c) \text{ or } (b + c) \cdot a = (b \cdot a) + (c \cdot a)$$

**Integral domain :** A ring is called an integral domain if :

- i. It is commutative
- ii. It has unit element
- iii. It is without zero divisors

**Field :** A ring  $R$  with at least two elements is called a field if it has following properties :

- i.  $R$  is commutative
  - ii.  $R$  has unity
  - iii.  $R$  is such that each non-zero element possesses multiplicative inverse.
- For example, the rings of real numbers and complex numbers are also fields.

**Que 2.31.** Consider a ring  $(R, +, \cdot)$  defined by  $a \cdot a = a$ , determine whether the ring is commutative or not.

AKTU 2014-15, Marks 05

**Answer**

Let  $a, b \in R$   $(a + b)^2 = (a + b)$

$$\Rightarrow (a + b)(a + b) = (a + b)$$

$$(a + b)a + (a + b)b = (a + b)$$

$$(a^2 + ba) + (ab + b^2) = (a + b)$$

$$(a + ba) + (ab + b) = (a + b)$$

$$(\because a^2 = a \text{ and } b^2 = b)$$

$$(a + b) + (ba + ab) = (a + b) + 0$$

$$\Rightarrow ba + ab = 0$$

$$a + b = 0 \Rightarrow a + b = a + a \text{ [being every element of its own additive inverse]}$$

$$\Rightarrow b = a$$

$$\Rightarrow ab = ba$$

$\therefore R$  is commutative ring.

**Que 2.32.** Write out the operation table for  $[Z_2, +_2, \cdot_2]$ . Is  $Z_2$  a ring? Is an integral domain? Is a field? Explain.

**Answer**

The operation tables are as follows :

we have  $Z_2 = \{0, 1\}$

$+_2$	0	1
0	0	1
1	1	0

$\cdot_2$	0	1
0	0	0
1	0	1

Since  $(Z_2, +_2, \cdot_2)$  satisfies the following properties :

- i. **Closure axiom :** All the entries in both the tables belong to  $Z_2$ . Hence, closure is satisfied.
- ii. **Commutative :** In both the tables all the entries about the main diagonal are same therefore commutativity is satisfied.
- iii. **Associative law :** The associative law for addition and multiplication are also satisfied.

- iv. Here 0 is the additive identity and 1 is the multiplicative identity. Identity property is satisfied.
- v. Inverse exists in both the tables. The additive inverse of 0, 1 are 1, 0 respectively and the multiplicative inverse of non-zero element of  $Z_2$  is 1.
- vi. Multiplication is distributive over addition.

Hence  $(Z_2, +_2, *_2)$  is a ring as well as field. Since, we know that every field is an integral domain therefore it is also an integral domain.

**Que 2.33.** If the permutation of the elements of {1, 2, 3, 4, 5} are given by  $a = (1\ 2\ 3)(4\ 5)$ ,  $b = (1)(2)(3)(4\ 5)$ ,  $c = (1\ 5\ 2\ 4)(3)$ . Find the value of  $x$ , if  $ax = b$ . And also prove that the set  $Z_4 = \{0, 1, 2, 3\}$  is a commutative ring with respect to the binary modulo operation  $+_4$  and  $*_4$ .

AKTU 2015-16, Marks 10

**Answer**

$$ax = b \Rightarrow (123)(45)x = (1)(2)(3)(4, 5)$$

$$\Rightarrow \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 3 & 1 & 4 & 5 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 5 & 4 \end{pmatrix} x = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 5 & 4 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 3 & 1 & 5 & 4 \end{pmatrix} x = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 5 & 4 \end{pmatrix}$$

$$x = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 3 & 1 & 5 & 4 \end{pmatrix}^{-1} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 5 & 4 \end{pmatrix}$$

$$= \begin{pmatrix} 2 & 3 & 1 & 5 & 4 \\ 1 & 2 & 3 & 4 & 5 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 5 & 4 \end{pmatrix}$$

$$= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 1 & 2 & 5 & 4 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 5 & 4 \end{pmatrix}$$

$$x = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 1 & 2 & 4 & 5 \end{pmatrix}$$

$+_4$	0	1	2	3
0	0	1	2	3
1	1	2	3	0
2	2	3	0	1
3	3	0	1	2

$*_4$	0	1	2	3
0	0	0	0	0
1	0	1	2	3
2	0	2	0	2
3	0	3	2	1

We find from these tables :

- i. All the entries in both the tables belong to  $Z_4$ . Hence,  $Z_4$  is closed with respect to both operations.

- ii. **Commutative law** : The entries of 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup> rows are identical with the corresponding elements of the 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup> columns respectively in both the tables. Hence,  $Z_4$  is commutative with respect to both operations.
- iii. **Associative law** : The associative law for addition and multiplication  $a +_4 (b +_4 c) = (a +_4 b) +_4 c$  for all  $a, b, c \in Z_4$   
 $a \times_4 (b \times_4 c) = (a \times_4 b) \times_4 c$ , for all  $a, b, c \in Z_4$   
 can easily be verified.
- iv. **Existence of identity** : 0 is the additive identity and 1 is multiplicative identity for  $Z_4$ .
- v. **Existence of inverse** : The additive inverses of 0, 1, 2, 3 are 0, 3, 2, 1 respectively.  
 Multiplicative inverse of non-zero element 1, 2, 3 are 1, 2, 3 respectively.
- vi. **Distributive law** : Multiplication is distributive over addition i.e.,

$$a \times_4 (b +_4 c) = a \times_4 b + a \times_4 c$$

$$(b +_4 c) \times_4 a = b \times_4 a + c \times_4 a$$

For,  $a \times_4 (b +_4 c) = a \times_4 (b + c)$  for  $b +_4 c = b + c \pmod{4}$

= least positive remainder when  $a \times (b + c)$  is divided by 4

= least positive remainder when  $ab + ac$  is divided by 4

$$= ab +_4 ac$$

$$= a \times_4 b +_4 a \times_4 c$$

For  $a \times_4 b = a \times b \pmod{4}$

Since  $(Z_4, +_4)$  is an abelian group,  $(Z_4, \times_4)$  is a semigroup and the operation is distributive over addition. The  $(Z_4, +_4, \times_4)$  is a ring. Now  $(Z_4, \times_4)$  is commutative with respect to  $\times_4$ . Therefore, it is a commutative ring.

**Que 2.34.** What is meant by ring? Give examples of both commutative and non-commutative rings.

AKTU 2018-19, Marks 07

**Answer**

Ring : Refer Q. 2.30, Page 2-19C, Unit-2.

Example of commutative ring : Refer Q. 2.31, Page 2-20C, Unit-2.

Example of non-commutative ring : Consider the set  $R$  of  $2 \times 2$  matrix with real element. For  $A, B, C \in R$

$$A * (B + C) = (A * B) + (A * C)$$

$$(A + B) * C = (A * C) + (B * C)$$

also,

$\therefore *$  is distributive over  $+$ .

$\therefore (R, +, *)$  is a ring.

We know that  $AB \neq BA$ , Hence  $(R, +, *)$  is non-commutative ring.

### VERY IMPORTANT QUESTIONS

*Following questions are very important. These questions may be asked in your SESSIONALS as well as UNIVERSITY EXAMINATION.*

**Q. 1.** Let  $H$  be a subgroup of a finite group  $G$ . Prove that order of  $H$  is a divisor of order of  $G$ .

**ANS:** Refer Q. 2.5.

**Q. 2.** Prove that  $(Z_6, (+_6))$  is an abelian group of order 6, where  $Z_6 = \{0, 1, 2, 3, 4, 5\}$ .

**ANS:** Refer Q. 2.7.

**Q. 3.** Let  $G = \{1, -1, i, -i\}$  with the binary operation multiplication be an algebraic structure, where  $i = \sqrt{-1}$ . Determine whether  $G$  is an abelian or not.

**ANS:** Refer Q. 2.8.

**Q. 4.** Write the properties of group. Show that the set  $\{1, 2, 3, 4, 5\}$  is not group under addition and multiplication modulo 6.

**ANS:** Refer Q. 2.9.

**Q. 5.** Let  $G$  be a group and let  $a, b \in G$  be any elements. Then

i.  $(a^{-1})^{-1} = a$

ii.  $(a * b)^{-1} = b^{-1} * a^{-1}$ .

**ANS:** Refer Q. 2.11.

**Q. 6.** Prove that the intersection of two subgroups of a group is also subgroup.

**ANS:** Refer Q. 2.12.

**Q. 7.** Let  $G$  be the set of all non-zero real number and let  $a * b = ab/2$ . Show that  $(G, *)$  be an abelian group.

**ANS:** Refer Q. 2.13.

**Q. 8.** Prove that inverse of each element in a group is unique.

**ANS:** Refer Q. 2.14.

**Q. 9.** Prove that every group of prime order is cyclic.

**ANS:** Refer Q. 2.16.

Q. 10. Show that every group of order 3 is cyclic.

ANS: Refer Q. 2.17.

Q. 11. Show that  $G = [(1, 2, 4, 5, 7, 8), X_9]$  is cyclic. How many generators are there? What are they?

ANS: Refer Q. 2.19.

Q. 12. State and prove Lagrange's theorem for group. Is the converse true?

ANS: Refer Q. 2.21.

Q. 13.

a. Prove that every cyclic group is an abelian group.

b. Obtain all distinct left cosets of  $\{(0), (3)\}$  in the group  $(\mathbb{Z}_6, +_6)$  and find their union.

c. Find the left cosets of  $\{(0), [3]\}$  in the group  $(\mathbb{Z}_6, +_6)$ .

ANS: Refer Q. 2.23.

Q. 14. Write and prove the Lagrange's theorem. If a group  $G = \{\dots, -3, 2, -1, 0, 1, 2, 3, \dots\}$  having the addition as binary operation. If  $H$  is a subgroup of group  $G$  where  $x^2 \in H$  such that  $x \in G$ . What is  $H$  and its left coset w.r.t 1?

ANS: Refer Q. 2.24.

Q. 15. Consider a ring  $(R, +, \cdot)$  defined by  $a \cdot a = a$ , determine whether the ring is commutative or not.

ANS: Refer Q. 2.31.

Q. 16. If the permutation of the elements of  $\{1, 2, 3, 4, 5\}$  are given by  $a = (1\ 2\ 3)(4\ 5)$ ,  $b = (1)(2)(3)(4\ 5)$ ,  $c = (1\ 5\ 2\ 4)(3)$ . Find the value of  $x$ , if  $ax = b$ . And also prove that the set  $\mathbb{Z}_4 = \{0, 1, 2, 3\}$  is a commutative ring with respect to the binary modulo operation  $+_4$  and  $\cdot_4$ .

ANS: Refer Q. 2.33.

Q. 17. What is meant by ring? Give examples of both commutative and non-commutative rings.

ANS: Refer Q. 2.34.



## 3

UNIT

Lattices and  
Boolean Algebra

## CONTENTS

<b>Part-1</b> :	Lattices : Definition, Properties of Lattices-Bounded, Complemented, Modular and Complete Lattice	3-2C to 3-10C
<b>Part-2</b> :	Boolean Algebra : Introduction, Axioms and Theorems of Boolean Algebra	3-10C to 3-13C
<b>Part-3</b> :	Algebraic Manipulation of Boolean Expressions, Simplification of Boolean Functions, Karnaugh Maps	3-13C to 3-17C
<b>Part-4</b> :	Logic Gates, Digital Circuits and Boolean Algebra	3-17C to 3-20C

3-1 C (CS/IT-Sem-3) Downloaded from TwoWaits

3-2 C (CS/IT-Sem-3)

Lattices and Boolean Algebra

## PART-1

*Lattices : Definition, Properties of Lattices-Bounded,  
Complemented, Modular and Complete Lattice.*

## Questions-Answers

Long Answer Type and Medium Answer Type Questions

**Que 3.1.** Define lattice. Give its properties.

## Answer

A lattice is a poset  $(L, \leq)$  in which every subset  $\{a, b\}$  consisting of 2 elements has least upper bound (*lub*) and greatest lower bound (*glb*). Least upper bound is known as join of  $a$  and  $b$ . Greatest

## PART-1

*Lattices : Definition, Properties of Lattices-Bounded, Complemented, Modular and Complete Lattice.*

## Questions-Answers

## Long Answer Type and Medium Answer Type Questions

**Que 3.1.** Define lattice. Give its properties.

**Answer**

A lattice is a poset  $(L, \leq)$  in which every subset  $\{a, b\}$  consisting of 2 elements has least upper bound (*lub*) and greatest lower bound (*glb*). Least upper bound of  $\{a, b\}$  is denoted by  $a \vee b$  and is known as join of  $a$  and  $b$ . Greatest lower bound of  $\{a, b\}$  is denoted by  $a \wedge b$  and is known as meet of  $a$  and  $b$ . Lattice is generally denoted by  $(L, \wedge, \vee)$ .

**Properties :**

Let  $L$  be a lattice and  $a, b \in L$  then

**1. Idempotent property :**

$$\text{i. } a \vee a = a \qquad \text{ii. } a \wedge a = a$$

**2. Commutative property :**

$$\text{i. } a \vee b = b \vee a \qquad \text{ii. } a \wedge b = b \wedge a$$

**3. Associative property :**

$$\text{i. } a \vee (b \vee c) = (a \vee b) \vee c \qquad \text{ii. } a \wedge (b \wedge c) = (a \wedge b) \wedge c$$

**4. Absorption property :**

$$\text{i. } a \vee (a \wedge b) = a \qquad \text{ii. } a \wedge (a \vee b) = a$$

$$\text{5. i. } a \vee b = b \text{ iff } a \preceq b \qquad \text{ii. } a \wedge b = a \text{ iff } a \preceq b$$

$$\text{iii. } a \wedge b = a \text{ iff } a \vee b = b$$

**6. Distributive Inequality :**

$$\text{i. } a \wedge (b \vee c) \geq (a \wedge b) \vee (a \wedge c) \qquad \text{ii. } a \vee (b \wedge c) \leq (a \vee b) \wedge (a \vee c)$$

**7. Isotonicity :**

Let  $a, b, c, \in L$  and  $a \preceq b$ , then

$$\text{i. } a \vee c \preceq b \vee c \qquad \text{ii. } a \wedge c \preceq b \wedge c$$

**8. Stability :**

Let  $a, b, c, \in L$ , then

$$\text{i. } a \preceq b, a \preceq c \Rightarrow a \preceq (b \vee c), a \preceq (b \wedge c)$$

$$\text{ii. } a \succeq b, a \succeq c \Rightarrow a \succeq (b \wedge c), a \succeq (b \vee c)$$

**Que 3.2.** Explain types of lattice.

**8. Stability :**

Let  $a, b, c, \in L$ , then

i.  $a \leq b, a \leq c \Rightarrow a \leq (b \vee c), a \leq (b \wedge c)$

ii.  $a \geq b, a \geq c \Rightarrow a \geq (b \wedge c), a \geq (b \vee c)$

**Que 3.2.** Explain types of lattice.

**Answer**

**Types of lattice :**

1. **Bounded lattice :** A lattice  $L$  is said to be bounded if it has a greatest element 1 and a least element 0. In such lattice we have

$$a \vee 1 = 1, a \wedge 1 = a$$

$$a \vee 0 = a, a \wedge 0 = 0$$

$$\forall a \in L \text{ and } 0 \leq a \leq 1$$

2. **Complemented lattice :** Let  $L$  be a bounded lattice with greatest element 1 and least element 0. Let  $a \in L$  then an element  $a' \in L$  is complement of  $a$  if,

$$a \vee a' = 1 \text{ and } a \wedge a' = 0$$

A lattice  $L$  is called complemented if is bounded and if every element in  $L$  has a complement.

3. **Distributive lattice :** A lattice  $L$  is said to be distributive if for any element  $a, b$  and  $c$  of  $L$  following properties are satisfied :

i.  $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$

ii.  $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$

otherwise  $L$  is non-distributive lattice.

4. **Complete lattice :** A lattice  $L$  is called complete if each of its non-empty subsets has a least upper bound and greatest lower bound.

**For example :**

i.  $(\mathbb{Z}, \leq)$  is not a complete lattice.

ii. Let  $S$  be the class of all subsets of some universal set  $A$  and a relation  $\leq$  is defined as  $X \leq Y \Rightarrow X$  is a subset of  $Y$  such that  $X \wedge Y = X \cap Y$  and  $X \cup Y$ . Every subset of  $S$  has *glb* and *lub*. So,  $S$  is complete.

5. **Modular lattice :** A lattice  $(L, \leq)$  is called modular lattice if,  $a \vee (b \wedge c) = (a \vee b) \wedge c$  whenever  $a \leq c$  for all  $a, b, c \in L$ .

**Que 3.3.** If the lattice is represented by the Hasse diagram given below :

i. Find all the complements of 'e'.

ii. Prove that the given lattice is bounded complemented lattice.




Fig. 3.3.1.

AKTU 2014-15, Marks 10



PhonePe - UPI Payments, Recharges & Money Transfer

by DemandScale



**Answer**

- i. Complements of  $e$  are  $c$  and  $d$  which are as follows :  
 $c \vee e = b$  ,  $c \wedge e = f$   
 $d \vee e = b$  ,  $d \wedge e = f$
- ii. A lattice is bounded if it has greatest and least elements. Here  $b$  is greatest and  $f$  is least element.

**Que 3.4.** Define a lattice. For any  $a, b, c, d$  in a lattice  $(A, \leq)$  if  $a \leq b$  and  $c \leq d$  then show that  $a \vee c \leq b \vee d$  and  $a \wedge c \leq b \wedge d$ .

AKTU 2018-19, Marks 07

**Answer**

Lattice : Refer Q. 3.1, Page 3-2C, Unit-3.

**Numerical :**

As  $a \leq b$  and  $c \leq d$ ,  $a \leq b \leq b \vee d$  and  $c \leq d \leq b \vee d$ .

By transitivity of  $\leq$ ,  $a \leq b \vee d$  and  $c \leq b \vee d$ .

So  $b \vee d$  is an upper bound of  $a$  and  $c$ .

So  $a \vee c \leq b \vee d$ .

As  $a \wedge c \leq a$  and  $a \wedge c \leq c$ ,  $a \wedge c \leq a \leq b$  and  $a \wedge c \leq c \leq d$ .

Hence  $a \wedge c$  is a lower bound of  $b$  and  $d$ . So  $a \wedge c \leq b \wedge d$ .

So  $a \wedge c \leq b \wedge d$ .

**Que 3.5.** Let  $L$  be a bounded distributed lattice, prove if a complement exists, it is unique. Is  $D_{12}$  a complemented lattice? Draw the Hasse diagram of  $[P(a, b, c), \subseteq]$ , (Note: ' $\subseteq$ ' stands for subset). Find greatest element, least element, minimal element and maximal element.

AKTU 2015-16, Marks 15

OR

Draw the Hasse diagram of  $[P(a, b, c), \subseteq]$  (Note: ' $\subseteq$ ' stands for subset). Find greatest element, least element, minimal element and maximal element.

AKTU 2015-16, Marks 10

**Answer**

Let  $a_1$  and  $a_2$  be two complements of an element  $a \in L$ .

Then by definition of complement

$$\left. \begin{aligned} a \vee a_1 &= I \\ a \wedge a_1 &= 0 \end{aligned} \right\} \dots(3.5.1)$$

$$\left. \begin{aligned} a \vee a_2 &= I \\ a \wedge a_2 &= 0 \end{aligned} \right\} \dots(3.5.2)$$

$$\left. \begin{aligned} a \vee a_2 = I \\ a \wedge a_2 = 0 \end{aligned} \right\}$$

...(3.5.2)

Consider

$$\begin{aligned} a_1 &= a_1 \vee 0 \\ &= a_1 \vee (a \wedge a_2) && \text{[from (3.5.2)]} \\ &= (a_1 \vee a) \wedge (a_1 \vee a_2) && \text{[Distributive property]} \\ &= (a \vee a_1) \wedge (a_1 \vee a_2) && \text{[Commutative property]} \\ &= I \wedge (a_1 \vee a_2) && \text{[from (3.5.1)]} \\ &= a_1 \vee a_2 && \dots(3.5.3) \end{aligned}$$

Now Consider

$$\begin{aligned} a_2 &= a_2 \vee 0 \\ &= a_2 \vee (a \wedge a_1) && \text{[from (3.5.2)]} \\ &= (a_2 \vee a) \wedge (a_2 \vee a_1) && \text{[Distributive property]} \\ &= (a \vee a_2) \wedge (a_1 \vee a_2) && \text{[Commutative property]} \\ &= I \wedge (a_1 \vee a_2) && \text{[from (3.5.1)]} \\ &= a_1 \vee a_2 && \dots(3.5.4) \end{aligned}$$

Hence, from (3.5.3) and (3.5.4),

$$a_1 = a_2$$

So, for bounded distributive lattice complement is unique.

Hasse diagram of  $[P(a, b, c), \subseteq]$  or  $[P(a, b, c), \leq]$  is shown in Fig. 3.5.1.




Fig. 3.5.1.

Greatest element is  $\{a, b, c\}$  and maximal element is  $\{a, b, c\}$ .  
The least element is  $\phi$  and minimal element is  $\phi$ .

**Que 3.6.** | The directed graph  $G$  for a relation  $R$  on set  $A = \{1, 2, 3, 4\}$  is shown below :




Fig. 3.6.1.

- Verify that  $(A, R)$  is a poset and find its Hasse diagram.
- Is this a lattice?
- How many more edges are needed in the Fig. 3.6.1 to extend  $(A, R)$  to a total order?
- What are the maximal and minimal elements?

AKTU 2014-15, Marks 10

Answer

- i. The relation  $R$  corresponding to the given directed graph is,  
 $R = \{(1, 1), (2, 2), (3, 3), (4, 4), (3, 1), (3, 4), (1, 4), (3, 2)\}$   
 $R$  is a partial order relation if it is reflexive, antisymmetric and transitive.

**Reflexive:** Since  $aRa, \forall a \in A$ . Hence, it is reflexive.

**Antisymmetric:** Since  $aRb$  and  $bRa$  then we get  $a = b$  otherwise  $aRb$  or  $bRa$ .

Hence, it is antisymmetric.

**Transitive:** For every  $aRb$  and  $bRc$  we get  $aRc$ . Hence, it is transitive.

Therefore, we can say that  $(A, R)$  is poset. Its Hasse diagram is :



Fig. 3.6.2.

- ii. Since there is no *lub* of 1 and 2 and same for 2 and 4. The given poset is not a lattice.

$\vee$	1	2	3	4
1	1	-	1	1
2	-	2	2	-
3	1	2	3	1
4	1	-	1	4

- iii. Only one edge  $(4, 2)$  is included to make it total order.  
 iv. Maximals are  $(1, 2)$  and minimal are  $(3, 4)$ .

**Que 3.7.** In a lattice if  $a \leq b \leq c$ , then show that

a.  $a \vee b = b \wedge c$

b.  $(a \vee b) \vee (b \wedge c) = (a \vee b) \wedge (a \vee c) = b$

AKTU 2016-17, Marks 10

Answer

- a. Given :  $a \leq b \leq c$



Now  $a \vee b =$  least upper bound of  $a, b$   
 $=$  least {all upper bounds of  $a, b$ }  
 $=$  least  $\{b, c, \dots\}$  [using  $a \leq b \leq c$ ]  
 $= b$  ... (3.7.1)

and  $b \wedge c =$  greatest lower bound of  $b, c$   
 $=$  maximum {all lower bounds of  $b, c$ }  
 $=$  maximum  $\{b, a, \dots\}$  [using  $a \leq b \leq c$ ]  
 $= b$  ... (3.7.2)

Eq. (3.7.1) and (3.7.2) gives,  $a \vee b = b \wedge c$

b.  $(a \vee b) \vee (b \wedge c) \Rightarrow (a \vee b) \wedge (a \vee c) = b$

Consider,  $(a \vee b) \vee (b \wedge c)$   
 $= b \vee b$  [using  $a \leq b \leq c$  and definition of  $\vee$  and  $\wedge$ ]  
 $= b$  ... (3.7.3)

and  $(a \vee b) \wedge (a \vee c) = b \wedge c$   
 $= b$  ... (3.7.4)

From eq. (3.7.3) and (3.7.4),  $(a \vee b) \vee (b \wedge c) = (a \vee b) \wedge (a \vee c) = b$ .

**Que 3.8.**

- a. Prove that every finite subset of a lattice has an LUB and a GLB.
- b. Give an example of a lattice which is a modular but not a distributive.

**AKTU 2016-17, Marks 10**

**Answer**

- a.
  1. The theorem is true if the subset has 1 element, the element being its own *glb* and *lub*.
  2. It is also true if the subset has 2 elements.
  3. Suppose the theorem holds for all subsets containing 1, 2, ...,  $k$  elements, so that a subset  $a_1, a_2, \dots, a_k$  of  $L$  has a *glb* and a *lub*.
  4. If  $L$  contains more than  $k$  elements, consider the subset  $\{a_1, a_2, \dots, a_{k+1}\}$  of  $L$ .
  5. Let  $w = \text{lub}(a_1, a_2, \dots, a_k)$ .
  6. Let  $t = \text{lub}(w, a_{k+1})$ .
  7. If  $s$  is any upper bound of  $a_1, a_2, \dots, a_{k+1}$ , then  $s$  is  $\geq$  each of  $a_1, a_2, \dots, a_k$  and therefore  $s \geq w$ .
  8. Also,  $s \geq a_{k+1}$  and therefore  $s$  is an upper bound of  $w$  and  $a_{k+1}$ .
  9. Hence  $s \geq t$ .
  10. That is, since  $t \geq$  each  $a_i$ ,  $t$  is the *lub* of  $a_1, a_2, \dots, a_{k+1}$ .
  11. The theorem follows for the *lub* by finite induction.

12. If  $L$  is finite and contains  $m$  elements, the induction process stops when  $k + 1 = m$ .

b.

1. The diamond is modular, but not distributive.
2. Obviously the pentagon cannot be embedded in it.
3. The diamond is not distributive :  

$$y \vee (x \wedge z) = y (y \vee x) \wedge (y \vee z) = 1$$
4. The distributive lattices are closed under sublattices and every sublattice of a distributive lattice is itself a distributive lattice.
5. If the diamond can be embedded in a lattice, then that lattice has a non-distributive sublattice, hence it is not distributive.

**Que 3.9.** Show that the inclusion relation  $\subseteq$  is a partial ordering on the power set of a set  $S$ . Draw the Hasse diagram for inclusion on the set  $P(S)$ , where  $S = \{a, b, c, d\}$ . Also determine whether  $(P(S), \subseteq)$  is a lattice.

**AKTU 2018-19, Marks 07**

**Answer**


Show that the inclusion relation ( $\subseteq$ ) is a partial ordering on the power set of a set  $S$ .

**Reflexivity :**  $A \subseteq A$  whenever  $A$  is a subset of  $S$ .

**Antisymmetry :** If  $A$  and  $B$  are positive integers with  $A \subseteq B$  and  $B \subseteq A$ , then  $A = B$ .

**Transitivity :** If  $A \subseteq B$  and  $B \subseteq C$ , then  $A \subseteq C$ .

**Hasse diagram :**



**Fig. 3.9.1.**

$(P(S), \subseteq)$  is not a lattice because  $(\{a, b\}, \{b, d\})$  has no *lub* and *glb*.


**Que 3.10.** Explain modular lattice, distributive lattice and bounded lattice with example and diagram. **AKTU 2017-18, Marks 10**

**Answer**

**Modular distributive and bounded lattice :** Refer Q. 3.2, Page 3-2C, Unit-3.

**Example :**

Let consider a Hasse diagram :



**Fig. 3.10.1.**

**Modular lattice :**

$0 \leq a$  i.e., taking  $b = 0$

$b \vee (a \wedge c) = 0 \vee 0 = 0, a \wedge (b \vee c) = a \wedge c = 0$

**Distributive lattice :**

For a set  $S$ , the lattice  $P(S)$  is distributive, since union and intersection each satisfy the distributive property.

**Bounded lattice :** Since, the given lattice has 1 as greatest and 0 as least element so it is bounded lattice.


**Que 3.11.** Draw the Hasse diagram of  $(A, \leq)$ , where

$A = \{3, 4, 12, 24, 48, 72\}$  and relation  $\leq$  be such that  $a \leq b$  if  $a$  divides  $b$ .

**AKTU 2017-18, Marks 07**

**Answer**

Hasse diagram of  $(A, \leq)$  where  $A = \{3, 4, 12, 24, 48, 72\}$



**Fig. 3.11.1.**

**Que 3.12.** For any positive integer  $D36$ , then find whether  $(D36, |)$  is lattice or not ?


**AKTU 2017-18, Marks 07**

**Answer**

D36 = Divisor of 36 = {1, 2, 3, 4, 6, 9, 12, 18, 36}

Hasse diagram :

$$(1 \vee 3) = \{3, 6\}, (1 \vee 2) = \{2, 4\}, (2 \vee 6) = \{6, 18\}, (9 \vee 4) = \{\phi\}$$



Since,

$$9 \vee 4 = \{\phi\}$$

So, D36 is not a lattice.

**PART-2**

*Boolean Algebra : Introduction, Axioms and Theorems of Boolean Algebra.*

**Questions-Answers**

Long Answer Type and Medium Answer Type Questions

**Que 3.13.** What is Boolean algebra? Write the axioms of Boolean algebra. Also, describe the theorems of it.

**Answer**

A Boolean algebra is generally denoted by  $(B, +, \cdot, 0, 1)$  where  $(B, +, \cdot)$  is a lattice with binary operations '+' and ' $\cdot$ ' called the join and meet respectively and ' $'\cdot'$ ' is unary operation in  $B$ . The elements 0 and 1 are zero (least) and unit (greatest) elements of lattice  $(B, +, \cdot)$ .  $B$  is called a Boolean algebra if the following axioms are satisfied for all  $a, b, c$  in  $B$ .

**Axioms of Boolean algebra :**

If  $a, b, c \in B$ , then

1. Commutative laws :

a.  $a + b = b + a$

b.  $a \cdot b = b \cdot a$

2. Distributive laws :

a.  $a + (b \cdot c) = (a + b) \cdot (a + c)$

b.  $a \cdot (b + c) = (a \cdot b) + (a \cdot c)$

3. Identity laws :

a.  $a + 0 = a$

b.  $a \cdot 1 = a$

- Distributive laws :
- a.  $a + (b.c) = (a + b).(a + c)$
  - b.  $a.(b + c) = (a.b) + (a.c)$
- Identity laws :
- a.  $a + 0 = a$
  - b.  $a.1 = a$

Discrete Structures & Theory of Logic

3-11 C (CSIT-Sem-3)

4. Complement laws :

- a.  $a + a' = 1$
- b.  $a.a' = 0$

Basic theorems :

Let  $a, b, c \in B$ , then

1. Idempotent laws :

- a.  $a + a = a$
- b.  $a.a = a$

2. Boundedness (Dominance) laws :

- a.  $a + 1 = 1$
- b.  $a.0 = 0$

3. Absorption laws :

- a.  $a + (a.b) = a$
- b.  $a.(a + b) = a$

4. Associative laws :

- a.  $(a + b) + c = a + (b + c)$
- b.  $(a.b).c = a.(b.c)$

5. Uniqueness of complement :

$a + x = 1$  and  $a.x = 0$ , then  $x = a'$

6. Involution law :  $(a')' = a$

- 7. a.  $0' = 1$
- b.  $1' = 0$

8. De-Morgan's laws :

- a.  $(a + b)' = a'.b'$
- b.  $(a.b)' = a' + b'$

**Que 3.14.** Prove the following theorems :

- a. Absorption law : Prove that  $\forall a, b, \in B$ 
  - i.  $a.(a + b) = a$
  - ii.  $a + a.b = a$
- b. Idempotent law : Prove that  $\forall a \in B, a + a = a$  and  $a.a = a$ .
- c. De Morgan's law : Prove that  $\forall a, b, \in B$ 
  - i.  $(a + b)' = a'.b'$
  - ii.  $(a.b)' = a' + b'$
- d. Prove that  $0' = 1$  and  $1' = 0$ .

**Answer**

a. Absorption law :

i. To prove :  $a.(a + b) = a$

Let  $a.(a + b) = (a + 0).(a + b)$   
 $= a + 0.b$   
 $= a + b.0$   
 $= a + 0$   
 $= a$

by Identity law  
 by Distributive law  
 by Commutative law  
 by Dominance law  
 by Identity law

ii. To prove :  $a + a.b = a$

Let  $a + a.b = a.1 + a.b$   
 $= a.(1 + b)$   
 $= a.(b + 1)$   
 $= a.1$   
 $= a$

by Identity law  
 by Distributive law  
 by Commutative law  
 by Dominance law  
 by Identity law

## b. Idempotent law :

To prove :  $a + a = a$  and  $a.a = a$ 

Let	$a = a + 0$	by Identity law
	$= a + a.a'$	by Complement law
	$= (a + a).(a + a')$	by Distributive law
	$= (a + a).1$	by Complement law
	$= a + a$	by Identity law
Now let	$a = a.1$	by Identity law
	$= a.(a + a')$	by Complement law
	$= a.a + a.a'$	by Distributive law
	$= a.a + 0$	by Complement law
	$= a.a$	by Identity law

## c. De Morgan's law :

i. To prove :  $(a + b)' = a'.b'$ 

To prove the theorem we will show that

$$(a + b) + a'.b' = 1$$

Consider	$(a + b) + a'.b' = [(a + b) + a'].[(a + b) + b']$	by Distributive law
	$= [(b + a) + a'].[(a + b) + b']$	
		by Commutative law
	$= (b + (a + a')).[a + (b + b')]$	
		by Associative law
	$= (b + 1).(a + 1)$	by Complement law
	$= 1.1$	by Dominance law
	$= 1$	...(3.14.1)

Also consider	$(a + b).a'b' = a'b'.(a + b)$	by Commutative law
	$= a'b'.a + a'b'.b$	by Distributive law
	$= a.(a'b') + a'.(b'b)$	by Commutative law
	$= (a.a').b' + a'.(b.b')$	by Associative law
	$= 0.b' + a'.0$	by Complement law
	$= b'.0 + a'.0$	by Commutative law
	$= 0 + 0$	by Dominance law
	$= 0$	...(3.14.2)

From eq. (3.14.1) and (3.14.2), we get,  
 $a'b'$  is complement of  $(a + b)$  i.e.  $(a + b)' = a'b'$ .

ii. To prove :  $(a.b)' = a' + b'$ 

Follows from principle of duality, that is, interchange operations  $+$  and  $\cdot$  and interchange the elements  $0$  and  $1$ .

d. To prove :  $0' = 1$  and  $1' = 0$ .

	$0' = (a.a)'$	by Complement law
	$= a' + (a)'$	by De Morgan's law
	$= a' + a$	by Involution law
	$= a + a'$	by Commutative law
	$= 1$	by Complement law

Now	$(0)' = 1'$	
-----	-------------	--

$\Rightarrow$	$0 = 1'$	
---------------	----------	--

$\Rightarrow$	$1' = 0$	
---------------	----------	--

# Take action NOW!



Discrete Structures & Theory of Logic

3-13 C (CS/IT-Sem-3)

**Que 3.15.** Define Boolean algebra. Show that in a Boolean algebra meet and join operations are distributive to each other.

**Answer**

**Boolean algebra :** Refer Q. 3.10, Page 3-10C, Unit-3.

**Meet and join operations are distributive :**

1. Let  $L$  be a poset under an ordering  $\leq$ . Let  $a, b \in L$ .
2. We define :  
 $a \vee b$  (read " $a$  join  $b$ ") as the least upper bound of  $a$  and  $b$ , and  
 $a \wedge b$  (read " $a$  meet  $b$ ") as greatest lower bound of  $a$  and  $b$ .
3. Since the join and meet operation produce a unique result in all cases where they exist, we can consider them as binary operations on a set if they always exist.
4. A lattice is a poset  $L$  (under  $\leq$ ) in which every pair of elements has a *lub* and a *glb*.
5. Since a lattice  $L$  is an algebraic system with binary operations  $\wedge$  and  $\vee$ , it is denoted by  $[L, \vee, \wedge]$ .
6. Let us consider,
  - a.  $[P(A), \vee, \wedge]$  is a lattice for any set  $A$  and
  - b. The join operation is the set operation of union and the meet operation is the operation of intersection; that is,  $\vee = \cup$  and  $\wedge = \cap$ .
7. It can be shown that the commutative laws, associative laws, idempotent laws, and absorption laws are all true for any lattice.
8. An example of this is clearly  $[P(A); \cup, \cap]$ , since these laws hold in the algebra of sets.
9. This lattice is also distributive such that join is distributive over meet and meet is distributive over join.

## PART-3

*Algebraic Manipulation of Boolean Expressions, Simplification of Boolean Functions, Karnaugh Maps.*

### Questions-Answers

**Long Answer Type and Medium Answer Type Questions**

**Que 3.16.** Express each Boolean expression as SOP and then in its complete sum-of-products form.

- a.  $E = x(xy' + x'y + y'z)$
- b.  $E = z(x' + y) + y'$

**Answer**

a.  $E = x.x.y' + x.x'.y + x.y'.z = x.y' + x.y'.z = x.y'$   
 ( $\because x.x'.y = 0$  and  $x.y'$  is contained in  $x.y'.z$ )  
 Complete SOP form  $= x.y'(z + z')$  ( $\because z$  is missing)  
 $= x.y'.z + x.y'.z'$

b.  $E = z.(x' + y) + y' = x'.z + y.z + y'$   
 Then  $E = x'.z(y + y') + y.z.(x + x') + y'.(x + x')(z + z')$   
 (by Complement law)  
 $= x'.y.z + x'.y'.z + x.y.z + x'.y.z + x.y'.z + x.z'.y'$   
 $+ x'.y'.z + x'.y'.z'$   
 $= x'.y.z + x'.y'.z + x.y.z + x.y'.z + x.y'.z' + x'.y'.z'$

**Que 3.17.** Define a Boolean function of degree  $n$ . Simplify the following Boolean expression using Karnaugh maps  
 $xyz + xy'z + x'y'z + x'yz + x'y'z'$

**Answer**

**Boolean function of degree  $n$  :**

1. Let  $B = \{0, 1\}$ . Then  $B^n = \{(x_1, x_2, \dots, x_n) \mid x_i \in B \text{ for } 1 \leq i \leq n\}$  is the set of all possible  $n$ -tuples of 0s and 1s.
2. The variable  $x$  is called a Boolean variable if it assumes values only from  $B$ , that is, if its only possible values are 0 and 1.
3. A function from  $B^n$  to  $B$  is called a Boolean function of degree  $n$ .
4. For example, the function  $F(x, y) = xy$  from the set of ordered pairs of Boolean variables to the set  $\{0, 1\}$ , is a Boolean function of degree 2 with  $F(1, 1) = 1, F(1, 0) = 0, F(0, 1) = 0$  and  $F(0, 0) = 0$ .

**Numerical :** The Karnaugh map for the given function is :

$xyz + xy'z + x'y'z + x'yz + x'y'z'$

	$yz$	$y'z'$	$y'z$	$yz$	$yz'$
$x'$	1	1	1		
$x$		1	1		

Fig. 3.17.1.

Then the simplified expression is :  $z + x'y'$ .

**Que 3.18.** Simplify the following Boolean functions using three variable maps :

- a.  $F(x, y, z) = \Sigma(0, 1, 5, 7)$
- b.  $F(x, y, z) = \Sigma(1, 2, 3, 6, 7)$

**Answer**

a.  $f(x, y, z) = \Sigma(0, 1, 5, 7)$

		0	1
z	xy		
	00	0	1
	01	2	3
	11	6	7
	10	4	5

$f = \bar{x}\bar{y} + xz$

**Fig. 3.18.1.**

b.  $f(x, y, z) = \Sigma(1, 2, 3, 6, 7)$

		00	01	11	10
x	yz				
	0	0	1	3	2
	1	4	5	7	6

**Fig. 3.18.2.**

$f = \bar{x}z + y$

**Que 3.19.** Simplify the following Boolean function using K-map :

$F(x, y, z) = \Sigma(0, 2, 3, 7)$

**AKTU 2017-18, Marks 07**

**Answer**

		00	01	11	10
x	yz				
	0	1		1	1
	1			1	

$F = \bar{x}\bar{z} + yz$

**Que 3.20.** Simplify the following Boolean expressions using K-map :

- a.  $Y = ((AB)' + A' + AB)'$
- b.  $A'B'C'D' + A'B'C'D + A'B'CD + A'B'CD' = A'B'$

**AKTU 2015-16, Marks 10**

**Answer**

a.  $Y = ((AB)' + A' + AB)'$   
 $= ((AB)')' (A' + AB)'$   
 $= (AB) ((A')' (AB)')$



$$\begin{aligned}
 &= AB (A (A' + B')) \\
 &= AB (AA' + AB') \\
 &= AB(0 + AB') = AB AB' \\
 &= ABB' \\
 &= 0
 \end{aligned}$$

Here, we find that the expression is not in minterm. For getting minterm, we simplify and find that its value is already zero. Hence, no need to use K-map for further simplification.

b.  $A'B'C'D' + A'B'C'D + A'B'CD + A'B'CD' = A'B'$   
 $= A'B'C'D' + A'B'C'D + A'B'CD + A'B'CD'$




Fig. 3.20.1.

On simplification by K-map, we get A'B' corresponding to all the four one's.

**Que 3.21.** Find the Boolean algebra expression for the following system. AKTU 2016-17, Marks 7.5




Fig. 3.21.1.

**Answer**




Fig. 3.21.2.

**Que 3.22.** Find the Sum-Of-Products and Product-Of-sum expansion of the Boolean function  $F(x, y, z) = (x + y)z'$ .

AKTU 2018-19, Marks 07

**Answer**

$$F(x, y, z) = (x + y)z'$$

x	y	z	x+y	z'	(x+y)z'
1	1	1	1	0	0
1	1	0	1	1	1
1	0	1	1	0	0
1	0	0	1	1	1
0	1	1	1	0	0
0	1	0	1	1	1
0	0	1	0	0	0
0	0	0	0	1	0

**Sum-Of-Product :**

$$F(x, y, z) = xyz' + xy'z' + x'yz'$$

**Product-Of-Sum :**

$$F(x, y, z) = (x + y + z)(x + y' + z)(x' + y + z)(x' + y' + z)$$

### PART-4

*Logic Gates, Digital Circuits and Boolean Algebra.*

### Questions-Answers

Long Answer Type and Medium Answer Type Questions

**Que 3.23.** Consider the Boolean function.

- a.  $f(x_1, x_2, x_3, x_4) = x_1 + (x_2 \cdot (x_1' + x_4)) + x_3 \cdot (x_2' + x_4')$
- Simplify  $f$  algebraically
  - Draw the logic circuit of the  $f$  and the reduction of the  $f$ .
- b. Write the expressions  $E_1 = (x + x * y) + (x/y)$  and  $E_2 = x + ((x * y + y)/y)$ , into

3-18 C (CS/IT-Sem-3)

Lattices and Boolean Algebra

- i. Prefix notation
- ii. Postfix notation

AKTU 2014-15, Marks 10

**Answer**

a. i.  $f(x_1, x_2, x_3, x_4) = x_1 + (x_2.(x_1' + x_4) + x_3.(x_2' + x_4'))$

$$= x_1 + x_2.x_1' + x_2.x_4 + x_3.x_2' + x_3.x_4'$$

$$= x_1 + x_2 + x_2.x_4 + x_3.x_2' + x_3.x_4'$$

$$= x_1 + x_2.(1 + x_4) + x_3.x_2' + x_3.x_4'$$

$$= x_1 + x_2 + x_3.x_2' + x_3.x_4'$$

$$= x_1 + x_2 + x_3 + x_3.x_4'$$

$$= x_1 + x_2 + x_3.(1 + x_4')$$

$$= x_1 + x_2 + x_3$$

ii. Logic circuit :



Fig. 3.23.1.

Reduction of f:

$$f(x_1, x_2, x_3, x_4) = x_1 + (x_2.(x_1' + x_4) + x_3.(x_2' + x_4'))$$

$$= x_1 + (x_2.x_1' + x_2.x_4) + (x_3.x_2' + x_3.x_4')$$

$$= x_1 + x_2.x_1' + x_2.x_4 + x_3.x_2' + x_3.x_4'$$

b.  $E_1 = (x + x * y) + (x/y)$

Binary tree is :




Fig. 3.23.2.

Prefix : ++ x \* x y / x y  
 Postfix : x x y \* + x y / +  
 $E_2 = x + ((x * y + y)/y)$

Binary tree is :




Fig. 3.23.3.

Prefix : + x / + \* x y y y

Postfix : x x y \* y + y / +

**Que 3.24.** Construct circuits that produce the following output :

$$F(X, Y, Z) = (X + Y + Z) (\bar{X}\bar{Y}\bar{Z})$$

**Answer**

The logic network is shown below :




Fig. 3.24.1.


**Que 3.25.** Draw the logic network corresponding to the following Boolean expressions :

- i.  $xy + x\bar{y}$
- ii.  $XY'Z' + XY'Z + XY'$




**Answer**

i. The logic network is shown in Fig. 3.25.1.



**Fig. 3.25.1.**

ii. The logic network shown in Fig. 3.25.2.




**Fig. 3.25.2.**

**VERY IMPORTANT QUESTIONS**

*Following questions are very important. These questions may be asked in your SESSIONALS as well as UNIVERSITY EXAMINATION.*

**Q. 1. If the lattice is represented by the Hasse diagram given below :**

- i. Find all the complements of 'e'.**
- ii. Prove that the given lattice is bounded complemented lattice.**



**Fig. 1.**




Fig. 1.

Downloaded from Two

**ANS.** Refer Q. 3.3.

**Q. 2.** Define a lattice. For any  $a, b, c, d$  in a lattice  $(A, \leq)$  if  $a \leq b$  and  $c \leq d$  then show that  $a \vee c \leq b \vee d$  and  $a \wedge c \leq b \wedge d$ .

**ANS.** Refer Q. 3.4.

**Q. 3.** Let  $L$  be a bounded distributed lattice, prove if a complement exists, it is unique. Is  $D_{12}$  a complemented lattice? Draw the Hasse diagram of  $[P(a, b, c), \subseteq]$ , (Note: ' $\subseteq$ ' stands for subset). Find greatest element, least element, minimal element and maximal element.

**ANS.** Refer Q. 3.5.

**Q. 4.** The directed graph  $G$  for a relation  $R$  on set  $A = \{1, 2, 3, 4\}$  is shown below :




Fig. 2.

- Verify that  $(A, R)$  is a poset and find its Hasse diagram.
- Is this a lattice?
- How many more edges are needed in the Fig. 2 to extend  $(A, R)$  to a total order?
- What are the maximal and minimal elements?

**ANS.** Refer Q. 3.6.

**Q. 5.** In a lattice if  $a \leq b \leq c$ , then show that

- $a \vee b = b \wedge c$
- $(a \vee b) \vee (b \wedge c) = (a \vee b) \wedge (a \vee c) = b$

**ANS.** Refer Q. 3.7.

**Q. 6.**

- Prove that every finite subset of a lattice has an LUB and a GLB.
- Give an example of a lattice which is a modular but not a distributive.

**ANS.** Refer Q. 3.8.

**Q. 7.** Show that the inclusion relation  $\subseteq$  is a partial ordering on the power set of a set  $S$ . Draw the Hasse diagram for inclusion on the set  $P(S)$ , where  $S = \{a, b, c, d\}$ . Also determine whether  $(P(S), \subseteq)$  is a lattice.

**ANS.** Refer Q. 3.9.

Downloaded from Two



# 4

UNIT

## Propositional Logic and Predicate Logic

### CONTENTS

- Part-1** : Propositional Logic : ..... 4-2C to 4-6C  
Proposition, Well Formed  
Formula, Truth Tables
- Part-2** : Tautology, Satisfiability, ..... 4-6C to 4-14C  
Contradiction, Algebra of  
Proposition, Theory of Inference
- Part-3** : Predicate Logic : ..... 4-15C to 4-19C  
First Order Predicate,  
Well Formed Formula  
of Predicate, Quantifiers,  
Inference Theory of  
Predicate Logic

**PART-1**

*Propositional Logic : Proposition, Well Formed Formula,  
Truth Tables.*

**Questions-Answers**

**Long Answer Type and Medium Answer Type Questions**

**Que 4.1.** Define the term proposition. Also, explain compound proposition with example.

**Answer**

**Proposition :** Proposition is a statement which is either true or false but not both. It is a declarative statement. It is usually denoted by lower case letters  $p, q, r, s, t$  etc. They are called Boolean variable or logic variable.  
**For example :**

1. Dr. A.P.J. Abdul Kalam was Prime Minister of India.
2. Roses are red.
3. Delhi is in India

(1) proposition is false whereas (2) and (3) are true.

**Compound proposition :** A compound proposition is formed by composition of two or more propositions called components or sub-propositions.

**For example :**

1. Risabh is intelligent and he studies hard.
2. Sky is blue and clouds are white.

Here first statement contain two propositions "Risabh is intelligent" and "he studies hard" whereas second statement contain propositions "sky is blue" and "clouds are white". As both statements are formed using two propositions. So they are compound propositions.

**Que 4.2.** Discuss connectives in detail with truth tables.

**Answer**

1. The words or phrases used to form compound proposition are called connectives.
2. There are five basic connectives as shown in the Table 4.2.1.

Table. 4.2.1.

S.No.	Connective words	Name of connective	Symbol
1.	Not	Negation	$\sim$ or $\neg$ or $-$
2.	And	Conjunction	$\wedge$
3.	Or	Disjunction	$\vee$
4.	If-then	Implication	$\rightarrow$
5.	If and only if	Biconditional	$\leftrightarrow$

- i. **Negation** : If  $P$  is a proposition then negation of  $P$  is a proposition which is true when  $p$  is false and false when  $p$  is true. It is denoted by  $\sim p$  or  $\neg$  or  $p'$  or  $\bar{p}$ .

**Truth table :**

$p$	$\sim p$
$T$	$F$
$F$	$T$

- ii. **Conjunction** : If  $p$  and  $q$  are two propositions then conjunction of  $p$  and  $q$  is a proposition which is true when both  $p$  and  $q$  are true otherwise false. It is denoted by  $p \wedge q$ .

**Truth table :**

$p$	$q$	$p \wedge q$
$T$	$T$	$T$
$T$	$F$	$F$
$F$	$T$	$F$
$F$	$F$	$F$

- iii. **Disjunction** : If  $p$  and  $q$  be two propositions, then disjunction of  $p$  and  $q$  is a proposition which is true when either one of  $p$  or  $q$  or both are true and is false when both  $p$  and  $q$  are false and it is denoted by  $p \vee q$ .

**Truth table :**

$p$	$q$	$p \vee q$
$T$	$T$	$T$
$T$	$F$	$T$
$F$	$T$	$T$
$F$	$F$	$F$

**Que 4.3.** What do you mean by well formed formula ?

**Answer**

1. Well formed formula is defined as a mathematical expression represented in well defined form and uses parenthesis, braces and square brackets to avoid the ambiguity.
2. Logical statements are also represented in well defined form using parenthesis according to priority of operations. To generate well formed formula recursively, following rules are used :
  - i. An atomic statement  $P$  is a well formed formula.
  - ii. If  $P$  is well formed formula then  $\sim P$  is also a well formed.
  - iii. If  $P$  and  $Q$  are well formed formulae then  $(P \vee Q)$ ,  $(P \wedge Q)$ ,  $(P \rightarrow Q)$  and  $(P \leftrightarrow Q)$  are also well formed formulae.
  - iv. A statement consists of variables, parenthesis and connectives is recursively a well formed formula iff it can be obtained by applying the above three rules.

**For example :**

1.  $(P \rightarrow (P \vee Q))$
2.  $((P \leftrightarrow Q) \rightarrow R)$

**Que 4.4.** Write short notes on :

- i. Truth table
- ii. Logical equivalence

**Answer**

- i. **Truth table :** A truth table is a table that shows the truth value of a compound proposition for all possible cases.

$p$	$q$	$(p \wedge q)$
$T$	$T$	$T$
$T$	$F$	$F$
$F$	$T$	$F$
$F$	$F$	$F$

$p$	$q$	$(p \vee q)$
$T$	$T$	$T$
$T$	$F$	$T$
$F$	$T$	$T$
$F$	$F$	$F$

$p$	$\sim p$
$T$	$F$
$F$	$T$

- ii. **Logical equivalence :** If two propositions  $P(p, q, \dots)$  and  $Q(p, q, \dots)$  where  $p, q, \dots$  are propositional variables, have the same truth values in every possible case, the propositions are called logically equivalent or simply equivalent, and denoted by

$$P(p, q, \dots) = Q(p, q, \dots)$$

**Que 4.5.** Explain the following terms with suitable example :

- i. Conjunction
- ii. Disjunction
- iii. Conditional

iv. Converse

v. Contrapositive

AKTU 2014-15, Marks 10

OR

Define inverse.

Answer

- i. **Conjunction** : If  $p$  and  $q$  are two statements, then conjunction of  $p$  and  $q$  is the compound statement denoted by  $p \wedge q$  and read as " $p$  and  $q$ ". Its truth table is,

$p$	$q$	$p \wedge q$
$T$	$T$	$T$
$T$	$F$	$F$
$F$	$T$	$F$
$F$	$F$	$F$

**Example :** $p$  : Ram is healthy. $q$  : He has blue eyes. $p \wedge q$  : Ram is healthy and he has blue eyes.

- ii. **Disjunction** : If  $p$  and  $q$  are two statements, the disjunction of  $p$  and  $q$  is the compound statement denoted by  $p \vee q$  and it is read as " $p$  or  $q$ ". Its truth table is,

$p$	$q$	$p \vee q$
$T$	$T$	$T$
$T$	$F$	$T$
$F$	$T$	$T$
$F$	$F$	$F$

**Example :** $p$  : Ram will go to Delhi. $q$  : Ram will go to Calcutta. $p \vee q$  : Ram will go to Delhi or Calcutta.

- iii. **Conditional** : If  $p$  and  $q$  are propositions. The compound proposition if  $p$  then  $q$  denoted by  $p \Rightarrow q$  or  $p \rightarrow q$  and is called conditional proposition or implication. It is read as "If  $p$  then  $q$ " and its truth table is,

$p$	$q$	$p \Rightarrow q$
$T$	$T$	$T$
$T$	$F$	$F$
$F$	$T$	$T$
$F$	$F$	$T$

**Example :**

$p$  : Ram works hard.

$q$  : He will get good marks.

$p \rightarrow q$  : If Ram works hard then he will get good marks.

**For converse and contrapositive :**

Let  $p$  : It rains.

$q$  : The crops will grow.

- iv. **Converse :** If  $p \Rightarrow q$  is an implication then its converse is given by  $q \Rightarrow p$  states that  $S$  : If the crops grow, then there has been rain.
- v. **Contrapositive :** If  $p \Rightarrow q$  is an implication then its contrapositive is given by  $\sim q \Rightarrow \sim p$  states that,
- $t$  : If the crops do not grow then there has been no rain.

**Inverse :**

If  $p \Rightarrow q$  is implication the inverse of  $p \Rightarrow q$  is  $\sim p \Rightarrow \sim q$ .

Consider the statement

$p$  : It rains.

$q$  : The crops will grow

The implication  $p \Rightarrow q$  states that,

$r$  : If it rains then the crops will grow.

The inverse of the implication  $p \Rightarrow q$ , namely  $\sim p \Rightarrow \sim q$  states that.

$u$  : If it does not rain then the crops will not grow.

## PART-2

*Tautology, Satisfiability, Contradiction, Algebra of Proposition,  
Theory of Inference.*

### Questions-Answers

**Long Answer Type and Medium Answer Type Questions**

**Que 4.6.** Explain tautologies, contradictions, satisfiability and contingency.

**Answer**

1. **Tautology** : Tautology is defined as a compound proposition that is always true for all possible truth values of its propositional variables and it contains  $T$  in last column of its truth table.

Propositions like,

- i. The doctor is either male or female.
- ii. Either it is raining or not.

are always true and are tautologies.

2. **Contradiction** : Contradiction is defined as a compound proposition that is always false for all possible truth values of its propositional variables and it contains  $F$  in last column of its truth table.

Propositions like,

- i.  $x$  is even and  $x$  is odd number.
- ii. Tom is good boy and Tom is bad boy.

are always false and are contradiction.

3. **Contingency** : A proposition which is neither tautology nor contradiction is called contingency.

Here the last column of truth table contains both  $T$  and  $F$ .

4. **Satisfiability** :

A compound statement formula  $A (P_1, P_2, \dots, P_n)$  is said to be satisfiable, if it has the truth value  $T$  for at least one combination of truth value of  $P_1, P_2, \dots, P_n$ .

**Que 4.7.** Write short note on algebra of propositions.

**Answer**

Proposition satisfies various laws which are useful in simplifying complex expressions. These laws are listed as :

1. **Idempotent laws** :
  - a.  $p \vee p \equiv p$
  - b.  $p \wedge p \equiv p$
2. **Associative laws** :
  - a.  $(p \vee q) \vee r \equiv p \vee (q \vee r)$
  - b.  $(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$
3. **Commutative laws** :
  - a.  $p \vee q \equiv q \vee p$
  - b.  $p \wedge q \equiv q \wedge p$

4. Distributive laws :
- $p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$
  - $p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$
5. Identity laws :
- $p \vee F \equiv p$
  - $p \vee T \equiv T$
  - $p \wedge F \equiv F$
  - $p \wedge T \equiv p$
6. Complement laws :
- $p \vee \sim P \equiv T$
  - $p \wedge \sim p \equiv F$
  - $\sim T \equiv F$
  - $\sim F \equiv T$
7. Involution law :
- $\sim(\sim p) \equiv p$
8. De Morgan's laws :
- $\sim(p \vee q) \equiv \sim p \wedge \sim q$
  - $\sim(p \wedge q) \equiv \sim p \vee \sim q$
9. Absorption laws :
- $p \vee (p \wedge q) \equiv p$
  - $p \wedge (p \vee q) \equiv p$

These laws can easily be verified using truth table.

**Que 4.8.** Discuss theory of inference in propositional logic.

**Answer**

Rules of inference are the laws of logic which are used to reach the given conclusion without using truth table. Any conclusion which can be derived using these laws is called valid conclusion and hence the given argument is valid argument.

- Modus ponens (Law of detachment)** : By this rule if an implication  $p \rightarrow q$  is true and the premise  $p$  is true then we can always conclude that  $q$  is also true.

The argument is of the form :

$$\begin{array}{l} p \rightarrow q \\ p \\ \hline \therefore q \end{array}$$

- Modus tollens (Law of contraposition)** : By this rule if an implication  $p \rightarrow q$  is true and conclusion  $q$  is false then the premise  $p$  must be false.
- The argument is of the form :

$$\begin{array}{r} p \rightarrow q \\ \underline{-q} \\ \therefore -p \end{array}$$

3. **Hypothetical syllogism** : By this rule whenever the two implications  $p \rightarrow q$  and  $q \rightarrow r$  are true then the implication  $p \rightarrow r$  is also true.  
The argument is of the form :

$$\begin{array}{r} p \rightarrow q \\ \underline{q \rightarrow r} \\ \therefore p \rightarrow r \end{array}$$

4. **Disjunctive syllogism** : By this rule if the premises  $p \vee q$  and  $\neg q$  are true then  $p$  is true.  
The argument is of the form :

$$\begin{array}{r} p \vee q \\ \underline{\neg q} \\ \therefore p \end{array}$$

5. **Addition** : By this rule if  $p$  is true then  $p \vee q$  is true regardless the truth value of  $q$ .  
The argument is of the form :

$$\begin{array}{r} p \\ \hline \therefore p \vee q \end{array}$$

6. **Simplification** : By this rule if  $p \wedge q$  is true then  $p$  is true.  
The argument is of form :

$$\begin{array}{r} \underline{p \wedge q} \\ \therefore p \quad \text{or} \quad \therefore q \end{array}$$

7. **Conjunction** : By this rule if  $p$  and  $q$  are true then  $p \wedge q$  is true.  
The argument is of the form :

$$\begin{array}{r} p \\ \underline{q} \\ \therefore p \wedge q \end{array}$$

8. **Constructive dilemma** : By this rule if  $(p \rightarrow q) \wedge (r \rightarrow s)$  and  $p \vee r$  are true then  $q \vee s$  is true.  
The argument is of form :

$$\begin{array}{r} (p \rightarrow q) \wedge (r \rightarrow s) \\ \underline{p \vee r} \\ \therefore q \vee s \end{array}$$

9. **Destructive dilemma** : By this rule if  $(p \rightarrow q) \wedge (r \rightarrow s)$  and  $\neg q \wedge \neg s$  are true.  
The argument is of the form :

$$\frac{(p \rightarrow q) \wedge (r \rightarrow s) \quad \sim q \wedge s}{\therefore \sim p \wedge r}$$

10. **Absorption** : By this rule if  $p \rightarrow q$  is true then  $p \rightarrow (p \wedge q)$  is true. The argument is of the form :

$$\frac{p \rightarrow q}{\therefore p \rightarrow (p \wedge q)}$$

**Que 4.9.** What do you mean by valid argument? Are the following arguments valid? If valid, construct a formal proof; if not, explain why.

For students to do well in discrete structure course, it is necessary that they study hard. Students who do well in courses do not skip classes. Student who study hard do well in courses. Therefore students who do well in discrete structure course do not skip class.

**Answer**

**Valid arguments :**

- An argument  $P_1, P_2, \dots, P_n \vdash Q$  is said to be valid if  $Q$  is true whenever all the premises  $P_1, P_2, \dots, P_n$  are true.
- For example : Consider the argument :  $p \rightarrow q, q \vdash p$ .

$p$	$q$	$p \rightarrow q$
T	T	T
T	F	F
F	T	T
F	F	T

where  $P$  denotes the premise and  $C$  denotes the conclusion.

- From the truth table we can see in first and third rows both the premises  $q$  and  $p \rightarrow q$  are true, but the conclusion  $p$  is false in third row. Therefore, this is not a valid argument.
  - First and third rows are called critical rows.
  - This method to determine whether the conclusion logically follows from the given premises by constructing the relevant truth table is called truth table technique.
  - Also, we can say the argument  $P_1, P_2, \dots, P_n \vdash Q$  is valid if and only if the proposition  $P_1 \wedge P_2 \wedge \dots \wedge P_n \rightarrow Q$  is true or we can say if  $P_1 \wedge P_2 \wedge \dots \wedge P_n \rightarrow Q$  is a tautology.
- For example : Consider the argument  $p \rightarrow q, p \vdash q$ .

Then from the truth table :

$p$	$q$	$p \rightarrow q$	$p \wedge p \rightarrow q$	$p \wedge (p \rightarrow q) \rightarrow q$
$T$	$T$	$T$	$T$	$T$
$T$	$F$	$F$	$F$	$T$
$F$	$T$	$T$	$F$	$T$
$F$	$F$	$T$	$F$	$T$

$p \wedge (p \rightarrow q) \rightarrow q$  is a tautology since the last column contains  $T$  only.

$\therefore p \rightarrow q, p \vdash q$  is a valid argument.

### Numerical :

Let the propositional variables be :

$p \rightarrow$  Do well in the course.

$q \rightarrow$  They study hard.

$r \rightarrow$  Do not skip classes.

- For students to do well in discrete structure course, it is necessary that they study hard :  $p \rightarrow q$
- Students who do well in the courses do not skip classes :  $p \rightarrow r$
- Students who study hard do well in courses :  $q \rightarrow p$
- Therefore, students who do well in discrete structure course do not skip classes :  $p \rightarrow r$

Therefore, we have,

Given :  $p \rightarrow q$  (I)       $p \rightarrow r$  (II)       $q \rightarrow p$  (III)      Conclusion :  $p \rightarrow r$  (IV)

**Proof :** Taking III and II together we get

$q \rightarrow p, p \rightarrow r$  gives  $q \rightarrow r$  (Using hypothetical syllogism)  
V

Now taking I and V

$p \rightarrow q$  and  $q \rightarrow r$  we get  $p \rightarrow r$  (Using hypothetical syllogism)

Hence,  $p \rightarrow r$  is conclusion, so it is valid.

Yes, the statement is valid.

### Que 4.10.

- Show that  $((p \vee q) \wedge \sim (\sim p \wedge (\sim q \vee \sim r))) \vee (\sim p \wedge \sim q) \vee (\sim p \vee r)$  is a tautology without using truth table.
- Rewrite the following arguments using quantifiers, variables and predicate symbols :
  - All birds can fly.
  - Some men are genius.
  - Some numbers are not rational.
  - There is a student who likes mathematics but not geography.

AKTU 2014-15, Marks 10

OR

Show that  $((p \vee q) \wedge \sim (\sim p \wedge (\sim q \vee \sim r))) \vee (\sim p \wedge \sim q) \vee (\sim p \vee r)$  is a tautology without using truth table.

AKTU 2018-19, Marks 07

**Answer**

i. We have

$$((p \vee q) \wedge \sim(\sim p \wedge (\sim q \vee \sim r))) \vee (\sim p \wedge \sim q) \vee (\sim p \vee r)$$

$$\equiv ((p \vee q) \wedge \sim(\sim p \wedge \sim(q \wedge r))) \vee (\sim(p \vee q) \vee \sim(p \vee r))$$

(Using De Morgan's Law)

$$\equiv [(p \vee q) \wedge (p \vee (q \wedge r))] \vee \sim((p \vee q) \wedge (p \vee r))$$

$$\equiv [(p \vee q) \wedge (p \vee q) \wedge (p \wedge r)] \vee \sim((p \vee q) \wedge (p \vee r))$$

(Using Distributive Law)

$$\equiv [((p \vee q) \wedge (p \vee q)) \wedge (p \vee r)] \vee \sim((p \vee q) \wedge (p \vee r))$$

$$\equiv ((p \vee q) \wedge (p \vee r)) \vee \sim((p \vee q) \wedge (p \vee r))$$

$$\equiv x \vee \sim x \text{ where } x = (p \vee q) \wedge (p \wedge r)$$

$$\equiv T$$

ii. a.  $\forall x [B(x) \Rightarrow F(x)]$ b.  $\exists x [M(x) \wedge G(x)]$ c.  $\sim [\exists(x) (N(x) \wedge R(x))]$ d.  $\exists x [S(x) \wedge M(x) \wedge \sim G(x)]$ 

**Que 4.11.** "If the labour market is perfect then the wages of all persons in a particular employment will be equal. But it is always the case that wages for such persons are not equal therefore the labour market is not perfect". Test the validity of this argument

using truth table.

**AKTU 2014-15, Marks 10****Answer**Let  $p_1$  : The labour market is perfect. $p_2$  : Wages of all persons in a particular employment will be equal. $\sim p_2$  : Wages for such persons are not equal. $\sim p_1$  : The labour market is not perfect.The premises are  $p_1 \Rightarrow p_2$ ,  $\sim p_2$  and the conclusion is  $\sim p_1$ . The argument $p_1 \Rightarrow p_2$ ,  $\sim p_2 \Rightarrow \sim p_1$  is valid if  $((p_1 \Rightarrow p_2) \wedge \sim p_2) \Rightarrow \sim p_1$  is a tautology.

Its truth table is,

$p_1$	$p_2$	$\sim p_1$	$\sim p_2$	$p_1 \Rightarrow p_2$	$(p_1 \Rightarrow p_2) \wedge \sim p_2$	$(p_1 \Rightarrow p_2 \wedge \sim p_2) \Rightarrow \sim p_1$
T	T	F	F	T	F	T
T	F	F	T	F	F	T
F	T	T	F	T	F	T
F	F	T	T	T	T	T

Since  $((p_1 \Rightarrow p_2) \wedge \sim p_2) \Rightarrow \sim p_1$  is a tautology. Hence, this is valid argument.**Que 4.12.** What is a tautology, contradiction and contingency ?Show that  $(p \vee q) \vee (\sim p \vee r) \rightarrow (q \vee r)$  is a tautology, contradiction or contingency.**AKTU 2018-19, Marks 07**

**Answer**

**Tautology, contradiction and contingency :** Refer Q. 4.6, Page 4-7C, Unit-4.

**Proof :**  $((p \vee q) \vee (\sim p \vee r)) \rightarrow (q \vee r)$

$p$	$q$	$r$	$\sim P$	$(p \vee q)$ $= A$	$(\sim p \vee r)$ $= B$	$(A \vee B)$ $= C$	$(q \vee r)$ $= D$	$C \rightarrow D$
F	F	F	T	F	T	T	F	F
F	F	T	T	F	T	T	T	T
F	T	F	T	T	T	T	T	T
F	T	T	T	T	T	T	T	T
T	F	F	F	T	F	T	F	F
T	F	T	F	T	T	T	T	T
T	T	F	F	T	F	T	T	T
T	T	T	F	T	T	T	T	T

So,  $((p \vee q) \vee (\sim p \vee r)) \rightarrow (q \vee r)$  is contingency.

**Que 4.13.**

- Find a compound proposition involving the propositional variables  $p$ ,  $q$ ,  $r$  and  $s$  that is true when exactly three propositional variables are true and is false otherwise.
- Show that the hypothesis "It is not sunny this afternoon and it is colder than yesterday", "We will go swimming only if it is sunny", "If we do not go swimming, then we will take a canoe trip." and "If we take a canoe trip, then we will be home by sunset" lead to the conclusion "We will be home by sunset."

OR

Show that the premises "It is not sunny this afternoon and it is colder than yesterday," "We will go swimming only if it is sunny," "If we do not go swimming, then we will take a canoe trip." and "If we take a canoe trip, then we will be home by sunset" lead to the conclusion "We will be home by sunset."

**AKTU 2018-19, Marks 07**

**Answer**

- The compound proposition will be :  $(p \wedge q \wedge r) \Leftrightarrow s$
- Let  $p$  be the proposition "It is sunny this afternoon",  $q$  be the proposition "It is colder than yesterday",  $r$  be the proposition "We will go swimming",  $s$  be the proposition "We will take a canoe trip", and  $t$  be the proposition "We will be home by sunset".

Then the hypothesis becomes  $\neg p \wedge q$ ,  $r \rightarrow p$ ,  $\neg r \rightarrow s$ , and  $s \rightarrow t$ . The conclusion is simply  $t$ .

We construct an argument to show that our hypothesis lead to the conclusion as follows :

S.No.	Step	Reason
1.	$\neg p \wedge q$	Hypothesis
2.	$\neg p$	Simplification using step 1
3.	$r \rightarrow p$	Hypothesis
4.	$\neg r$	Modus tollens using steps 2 and 3
5.	$\neg r \rightarrow s$	Hypothesis
6.	$s$	Modus ponens using steps 4 and 5
7.	$s \rightarrow t$	Hypothesis
8.	$t$	Modus ponens using steps 6 and 7

**Que 4.14.** Show that :  $(r \rightarrow \sim q, r \vee S, S \rightarrow \sim q, p \rightarrow q) \leftrightarrow \sim p$  are inconsistent.

AKTU 2017-18, Marks 07

### Answer

Following the indirect method, we introduce  $p$  as an additional premise and show that this additional premise leads to a contradiction.

(1)	(1) $p \rightarrow q$	Rule P
(2)	(2) $p$	Rule P (assumed)
{1, 2}	(3) $q$	Rule T, (1), (2) and modus ponens
(4)	(4) $s \rightarrow \bar{q}$	Rule P
{1, 2, 4}	(5) $\bar{s}$	Rule T, (3), (4) and modus tollens
(6)	(6) $r \vee s$	Rule P
{1, 2, 4, 6}	(7) $r$	Rule T, (5), (6) disjunctive syllogism
(8)	(8) $r \rightarrow \bar{q}$	Rule P
(8)	(9) $\bar{r} \vee \bar{q}$	Rule T, (8) and $EQ_{16} (p \rightarrow q \equiv \bar{p} \vee q)$
(8)	(10) $\overline{r \wedge q}$	Rule T, (8) and De Morgan's law
{1, 2, 4, 6}	(11) $r \wedge q$	Rule T, (7), (3) and conjunction
{1, 2, 4, 6, 8}	(12) $r \wedge q \wedge \overline{r \wedge q}$	Rule T, (10), (11) and conjunction.

Since, we know that set of formula is inconsistent if their conjunction implies contradiction. Hence it leads to a contradiction. So, it is inconsistent.

Since, we know that set of formula is inconsistent if their conjunction in contradiction. Hence it leads to a contradiction. So, it is

**PART-3**

*Predicate Logic : First Order Predicate, Well Formed Formula of Predicate, Quantifiers, Inference Theory of Predicate Logic.*

**Questions-Answers**

**Long Answer Type and Medium Answer Type Questions**

**Que 4.15.** Write a short note on

1. First order logic
2. Quantifiers

**Answer**

1. **First order logic :**

- i. First order logic is the extension of propositional logic by generalizing and quantifying the propositions over given universe of discourse.
- ii. In first order logic every individual has the property  $p$  (say).
- iii. It is also called first order predicate calculus.
- iv. Predicate calculus is generalization of propositional calculus. Predicate calculus allows us to manipulate statements about all or something.
- v. Universe of Discourse (UD) : It is the set of all possible values that can be substituted in place of predicate variable.

2. **Quantifiers :** There are following two types of quantifiers :

i. **Universal quantifier :** Let  $p(x)$  be a propositional function defined on set  $A$ . Consider the expression.

$$(\forall x \in A) P(x) \text{ or } \forall x P(x) \quad \dots(4.15.1)$$

Here the symbol " $\forall$ " is read as "for all" or "for every" and is called universal quantifier. Then the statement (4.15.1) is read as "For every  $x \in A$ ,  $P(x)$  is true."

ii. **Existential quantifier :** Let  $Q(x)$  be a propositional function defined on set  $B$ . Consider the expression

$$(\exists x \in B) Q(x) \text{ or } \exists x Q(x) \quad \dots(4.15.2)$$

Here the symbol " $\exists$ " is read as "for some" or "for at least one" or "there exists" and is called existential quantifier.

Then the statement (4.15.2) is read as "For some  $x \in B$ ,  $Q(x)$  is true".

**Que 4.16.** Explain rules of inference in predicate logic.

**Answer**

Rule of inference is a logical form consisting of function which takes premises, analyzes their syntax and returns a conclusion.

**Rules of inference :**

- i. **Universal specification :** By this rule if the premise  $(\forall x) P(x)$  is true then  $P(c)$  is true where  $c$  is particular member of UD.

$$\frac{(\forall x)P(x)}{\therefore P(c)}$$

- ii. **Universal generalization :** By this rule if  $P(c)$  is true for all  $c$  in UD then  $(\forall x) P(x)$  is true.

$$\frac{P(c)}{\therefore (\forall x)P(x)}$$

$x$  is not free in any of given premises.

- iii. **Existential specification :** By this rule if  $(\exists x) P(x)$  is true then  $P(x)$  is true for some particular member of UD.

$$\frac{(\exists x)P(x)}{\therefore P(c)}$$

$c$  is some member of UD.

- iv. **Existential generalization :** By this rule if  $P(c)$  is true for some particular member  $c$  in UD, then  $(\exists x) P(x)$  is true

$$\frac{P(c)}{\therefore (\exists x)P(x)}$$

$c$  is some member of UD.

- v. **Universal modus ponens :** By this rule if  $P(x) \rightarrow Q(x)$  is true for every  $x$  and  $P(c)$  is true for some particular member  $c$  in UD then  $Q(c)$  is true.

$$\frac{(\forall x) P(x) \rightarrow Q(x) \quad P(c)}{\therefore Q(c)}$$

- vi. **Universal modus tollens :** By this rule if  $P(x) \rightarrow Q(x)$  is true for every  $x$  and  $\sim Q(c)$  is true for some particular  $c$  in UD then  $\sim P(c)$  is true.

$$\frac{(\forall x) P(x) \rightarrow Q(x) \quad \sim Q(c)}{\therefore \sim P(c)}$$

**Que 4.17.** Write the symbolic form and negate the following statements :

- Everyone who is healthy can do all kinds of work.
- Some people are not admired by everyone.
- Everyone should help his neighbours, or his neighbours will not help him.  $e$

AKTU 2016-17, Marks 10

**Answer****a. Symbolic form :**

Let  $P(x)$ :  $x$  is healthy and  $Q(x)$ :  $x$  do all work

$$\forall x(P(x) \rightarrow Q(x))$$

$$\text{Negation : } \neg(\forall x(P(x) \rightarrow Q(x)))$$

**b. Symbolic form :**

Let  $P(x)$  :  $x$  is a person

$A(x, y)$  :  $x$  admires  $y$

The given statement can be written as "There is a person who is not admired by some person" and it is  $(\exists x)(\exists y)[P(x) \wedge P(y) \wedge \neg A(x, y)]$

$$\text{Negation : } (\exists x)(\exists y)[P(x) \wedge P(y) \wedge A(x, y)]$$

**c. Symbolic form :**

Let  $N(x, y)$  :  $x$  and  $y$  are neighbours

$H(x, y)$  :  $x$  should help  $y$

$P(x, y)$  :  $x$  will help  $y$

The statement can be written as "For every person  $x$  and every person  $y$ , if  $x$  and  $y$  are neighbours, then either  $x$  should help  $y$  or  $y$  will not help  $x$ " and it is  $(\forall x)(\forall y)[N(x, y) \rightarrow (H(x, y) \vee \neg P(y, x))]$

$$\text{Negation : } (\exists x)(\exists y)[N(x, y) \wedge \neg(H(x, y) \vee \neg P(y, x))]$$

**Que 4.18.** Express the following statements using quantifiers and logical connectives.

- Mathematics book that is published in India has a blue cover.
- All animals are mortal. All human being are animal. Therefore, all human being are mortal.
- There exists a mathematics book with a cover that is not blue.
- He eats crackers only if he drinks milk.
- There are mathematics books that are published outside India.
- Not all books have bibliographies. **AKTU 2015-16, Marks 10**

**Answer**

a.  $P(x)$  :  $x$  is a mathematic book published in India

$Q(x)$  :  $x$  is a mathematic book of blue cover

$$\forall x P(x) \rightarrow Q(x).$$

b.  $P(x)$  :  $x$  is an animal

$Q(x)$  :  $x$  is mortal

$$\forall x P(x) \rightarrow Q(x)$$

$R(x)$  :  $x$  is a human being

$\forall x R(x) \rightarrow P(x)$ .

c.  $P(x)$  :  $x$  is a mathematics book

$Q(x)$  :  $x$  is not a blue color

$\exists x, P(x) \wedge Q(x)$ .

d.  $P(x)$  :  $x$  drinks milk

$Q(x)$  :  $x$  eats crackers

for  $x$ , if  $P(x)$  then  $Q(x)$ .

or  $x, P(x) \Rightarrow Q(x)$ .

e.  $P(x)$  :  $x$  is a mathematics book

$Q(x)$  :  $x$  is published outside India

$\exists x P(x) \wedge Q(x)$ .

f.  $P(x)$  :  $x$  is a book having bibliography -  $\forall x, P(x)$ .

#### Que 4.19.

- Express this statement using quantifiers : "Every student in this class has taken some course in every department in the school of mathematical sciences".
- If  $\forall x \exists y P(x, y)$  is true, does it necessarily follow that  $\forall y P(x, y)$  is true? Justify your answer.

#### Answer

i.  $\forall x p(x) \Rightarrow \exists \forall z Q(y, z)$

where  $P(x)$  is student of class.

$Q(y, z)$  is the course from department.

- Let  $P(x, y)$  be  $x + y = 3$  and  $x, y$  belong to some set of integers  $\forall x \exists y P(x, y)$  is true means for all  $x$  there exists some  $y$  for which  $x + y = 3$  is true but for all  $y$  we conclude that  $P(x, y)$  will not be true.

**Que 4.20.** Translate the following sentences in quantified expressions of predicate logic.

- All students need financial aid.
- Some cows are not white.
- Suresh will get if division if and only if he gets first div.
- If water is hot, then Shyam will swim in pool.
- All integers are either even or odd integer.

AKTU 2017-18, Marks 07

**Answer**

- i.  $\forall x [S(x) \Rightarrow F(x)]$   
 ii.  $\sim [\exists(x) (C(x) \wedge W(x))]$   
 iii. Sentence is incorrect so cannot be translated into quantified expression.  
 iv.  $W(x) : x$  is water  
 $H(x) : x$  is hot  
 $S(x) : x$  is Shyam  
 $P(x) : x$  will swim in pool  
 $\forall x [(W(x) \wedge H(x)) \Rightarrow (S(x) \wedge P(x))]$   
 v.  $E(x) : x$  is even  
 $O(x) : x$  is odd  
 $\forall x (E(x) \vee O(x))$

**VERY IMPORTANT QUESTIONS**

*Following questions are very important. These questions may be asked in your SESSIONALS as well as UNIVERSITY EXAMINATION.*

**Q. 1. Explain the following terms with suitable example :**

- i. Conjunction
- ii. Disjunction
- iii. Conditional
- iv. Converse
- v. Contrapositive

**ANS:** Refer Q. 4.5.

**Q. 2. Show that  $((p \vee q) \wedge \sim(\sim p \wedge (\sim q \vee \sim r))) \vee (\sim p \wedge \sim q) \vee (\sim p \vee r)$  is a tautology without using truth table.**

**ANS:** Refer Q. 4.10.

**Q. 3. "If the labour market is perfect then the wages of all persons in a particular employment will be equal. But it is always the case that wages for such persons are not equal therefore the labour market is not perfect". Test the validity of this argument using truth table.**

**ANS:** Refer Q. 4.11.

**Q. 4. What is a tautology, contradiction and contingency? Show that  $(p \vee q) \vee (\sim p \vee r) \rightarrow (q \vee r)$  is a tautology, contradiction or contingency.**

**ANS.** Refer Q. 4.12.

**Q. 5.** Show that the premises "It is not sunny this afternoon and it is colder than yesterday," "We will go swimming only if it is sunny," "If we do not go swimming, then we will take a canoe trip." and "If we take a canoe trip, then we will be home by sunset" lead to the conclusion "We will be home by sunset."

**ANS.** Refer Q. 4.13.

**Q. 6.** Show that :  $(r \rightarrow \sim q, r \vee S, S \rightarrow \sim q, p \rightarrow q) \leftrightarrow \sim p$  are inconsistent.

**ANS.** Refer Q. 4.14.

**Q. 7.** Write the symbolic form and negate the following statements :

- Everyone who is healthy can do all kinds of work.
- Some people are not admired by everyone.
- Everyone should help his neighbours, or his neighbours will not help him.

**ANS.** Refer Q. 4.17.

**Q. 8.** Express the following statements using quantifiers and logical connectives.

- Mathematics book that is published in India has a blue cover.
- All animals are mortal. All human being are animal. Therefore, all human being are mortal.
- There exists a mathematics book with a cover that is not blue.
- He eats crackers only if he drinks milk.
- There are mathematics books that are published outside India.
- Not all books have bibliographies.

**ANS.** Refer Q. 4.18.

**Q. 9.** Translate the following sentences in quantified expressions of predicate logic.

- All students need financial aid.
- Some cows are not white.
- Suresh will get first division if and only if he gets first div.
- If water is hot, then Shyam will swim in pool.
- All integers are either even or odd integer.

**ANS.** Refer Q. 4.20.



# 5 UNIT

## Trees, Graph and Combinatorics

### CONTENTS

- Part-1** : Trees : Definition, ..... 5-2C to 5-4C  
Binary Tree
- Part-2** : Binary Tree Traversal ..... 5-5C to 5-8C
- Part-3** : Binary Search Tree ..... 5-8C to 5-12C
- Part-4** : Graphs : Definition ..... 5-12C to 5-15C  
and Terminology  
Representation of Graph
- Part-5** : Multigraph Bipartite ..... 5-15C to 5-24C  
Graphs, Planar Graph  
Isomorphism and  
Homomorphism of Graphs
- Part-6** : Graph Coloring ..... 5-24C to 5-24C
- Part-7** : Recurrence Relation and ..... 5-24C to 5-36C  
Generating Function :  
Recursive Definition of  
Function, Recursive  
Algorithms, Method of Solving  
Recurrences, Combinatorics :  
Introduction, Counting Techniques,  
Pigeonhole Principle


**Que 5.1.** Explain the following terms :

- i. Tree
- ii. Forest
- iii. Binary tree
- iv. Complete binary tree
- v. Full binary tree

**Answer**

**i. Tree :**

1. A tree is a connected graph that contains no cycle or circuit. It is a simple graph having no self loop or parallel edges.
2. A tree contains a finite set of elements which are called vertices or nodes. The vertex can have minimum degree 1 and maximum degree  $n$ .
3. The number of vertices in a tree is called order of tree.
4. A tree with only one vertex is called trivial or degenerate or empty tree.



**Fig. 5.1.1.**

**ii. Forest :** A forest is a collection of disjoint tree. It is an undirected, disconnected, acyclic graph.



**Fig. 5.1.2.**

**iii. Binary tree :** Binary tree is the tree in which the degree of every node is less than or equal to 2. A tree consisting of no nodes is also a binary tree.

- iv. **Complete binary tree** : A binary tree is said to be complete binary tree if all its levels, except the last, have maximum number of possible nodes (i.e., 2), and if all the nodes at the last level appear as far left as possible.




Fig. 5.1.3.

It is represented as  $T_n$  where  $n$  is number of nodes.

- v. **Full binary tree** : A binary tree is said to be extended or full binary tree if each node has either 0 or 2 children.




Fig. 5.1.4.

**Que 5.2.** Consider the tree given below :




Fig. 5.2.1.

- i. Which node is root ?
- ii. Which nodes are leaves ?
- iii. Name parent node of each ?
- iv. List children of each node.

- v. List the siblings.  
 vi. Find depth of each node.  
 vii. Find level of each node.

**Answer**

- i. The node  $a$  is the root node.  
 ii. The node  $g, h, i, j$  and  $l$  are leaves.

iii.

Nodes	Parent node
b, c	a
d	b
j, k	c
e, f	d
g, h	e
i	f
l	k

iv.

Nodes	Children
a	b, c
b	d
c	j, k
d	e, f
e	g, h
f	i
k	l

- v.  $b$  and  $c$   
 $e$  and  $f$   
 $g$  and  $h$   
 $j$  and  $k$

vi.

Nodes	Depth
a	1
b, c	2
d, j, k	3
e, f, l	4
g, h, i	5

vii.

Nodes	Level
a	0
b, c	1
d, j, k	2
e, f, l	3
g, h, i	4

**PART-2***Binary Tree Traversal.***Questions-Answers****Long Answer Type and Medium Answer Type Questions**

**Que 5.3.** Define preorder, inorder, and postorder tree traversal. Give an example of preorder, postorder, and inorder traversal of a binary tree of your choice with at least 12 vertices.

OR

Explain in detail about the binary tree traversal with an example.


**AKTU 2016-17, Marks 10**

**Answer**

**Tree traversal :** A traversal of tree is a process in which each vertex is visited exactly once in a certain manner. For a binary tree we have three types of traversal :

1. **Preorder traversal :** Each vertex is visited in the following order :
  - a. Visit the root (N).
  - b. Visit the left child (or subtree) of root (L).
  - c. Visit the right child (or subtree) of root (R).
2. **Postorder traversal :**
  - a. Visit the left child (subtree) of root.
  - b. Visit the right child (subtree) of root.
  - c. Visit the root.
3. **Inorder traversal :**
  - a. Visit the left child (subtree) of root.
  - b. Visit the root.
  - c. Visit the right child (subtree) of root.

**A binary tree with 12 vertices :**



**Fig. 5.3.1.**

Preorder (NLR) :  $ABDHIEJCFKLG$

Inorder (LNR) :  $HDIBJEAKFLCG$

Postorder (LRN) :  $HIDJEBKLF GCA$

**Que 5.4.** Construct a tree whose inorder and preorder traversal are as follows :

Inorder :  $Q \quad B \quad A \quad G \quad C \quad P \quad E \quad D \quad R$

Preorder :  $G \quad B \quad Q \quad A \quad P \quad C \quad D \quad E \quad R$


**Answer**

Root is  $G$ .




**Fig. 5.4.1.**

Element to left of  $G$  in inorder traversal are  $Q B A$  and  $B$  comes first in preorder traversal. Therefore tree will be



**Fig. 5.4.2.**

Now elements on the right of  $G$  in inorder traversal are  $C P E D R$  and  $P$  comes first. Therefore tree will be



**Fig. 5.4.3.**

Now element to left of  $P$  in inorder traversal is  $C$  and to the right is  $EDR$  (right subtree) out of  $EDR$ ,  $D$  comes first and  $E$  and  $R$  on its left and right respectively.

Therefore, the complete binary will look like




Fig. 5.4.4.

**Que 5.5.** Define a binary tree. A binary tree has 11 nodes. It's

inorder and preorder traversals node sequences are :

Preorder : *ABDHIEJL CFG*

Inorder : *HDIBJEK A F C G*

Draw the tree.


AKTU 2018-19, Marks 07

**Answer**


Binary tree : Refer Q. 5.1(iii), Page 5-2C, Unit-5.

Numerical :

**Step 1 :** In preorder sequence, leftmost element is the root of the tree. By searching A in 'Inorder Sequence' we can find out all the elements on the left and right sides of 'A'.



**Step 2 :** We recursively follow the above steps and we get



**Que 5.6.** Given the inorder and postorder traversal of a tree T :

Inorder : *HFEABIGDC* Postorder : *BEHFACDGI*


Determine the tree T and it's Preorder.

AKTU 2017-18, Marks 07


**Answer**

The root of tree is *I*.


Now elements on right of *I* are *D, G, C* and *G* comes last of all in postorder traversal.




Now *D* and *C* are on right of *G* and *D* comes last of *G* and *I* postorder traversal so



Now element left of *I* are *HFEAB* in inorder traversal and *A* comes last of all in postorder traversal. Therefore tree will be



Now *HFE* are on left of *A* in inorder traversal and *B* comes last of all and continuing in same manner. We will get final binary tree as



Preorder traversal of above binary tree is *CAFHEBDGI*

### PART-3

#### Binary Search Tree.

#### Questions-Answers

#### Long Answer Type and Medium Answer Type Questions


**Que 5.7.** What is a binary search tree? Form a binary search tree for the words vireo, warbler, egret, grosbeak, nuthatch, and kingfisher. Explain each step.

Download



**Answer****Binary search tree :**


1. A binary search tree is a binary tree  $T$  in which data is associated with the vertices.
2. The data are arranged so that, for each vertex  $v$  in  $T$ , each data item in the left subtree of  $v$  is less than the data item in  $v$  and each data item in the right subtree of  $v$  is greater than the data item in  $v$ .
3. The binary tree  $T$  in Fig. 5.7.1, is a binary search tree since every vertex in  $T$  exceeds every number in its left subtree and is less than every number in its right subtree.

**Fig. 5.7.1. A binary search tree.****Numerical :**


1. To start, create a vertex and place the first item in the list in this vertex and assign this as the key of the root.

vireo

2. To add warbler, compare it with vireo. Warbler is greater than vireo, therefore it will be in the right of vireo. *i.e.*,



3. Similarly by doing the above steps for rest of words we get the final binary search tree that is given below :



**Que 5.8.** Write algorithm for following :

- i. Searching and inserting a node in BST
- ii. Deleting a node in BST

**Answer**

i. Searching and inserting a node in BST :

Consider a binary search tree  $T$  and we have to insert or search an  $ITEM$  in the tree.

**Step I :** Compare  $ITEM$  with the root of the tree

- a. If  $ITEM < \text{root}$ , proceed to the left child of  $N$ .
- b. If  $ITEM > \text{root}$ , proceed to the right child of  $N$ .

**Step II :** Repeat Step I and if

- a. We reach a node  $N$  such that  $ITEM = N$  then search is successful.
- b. We reach an empty subtree, which shows the search is successful.

Insert  $ITEM$  in place of empty subtree.

ii. Deleting a node in BST :

Consider a binary search tree  $T$  and we have to delete an  $ITEM$  from the tree.

**Step I :** Find the location of node which contain  $ITEM$  and also keep track of its parent too.

**Step II :** Find the number of children of the node.

- a. If node has no children then simply delete the node.
- b. If the node has exactly one child then the node is deleted from tree by replacing the node from its child.
- c. If the node has two children then
  - i. Find its inorder successor.
  - ii. Delete the successor from tree using (a) or (b).
  - iii. Replace node by its successor in tree.


**Que 5.9.** Draw a binary search tree by inserting following integers 55, 20, 63, 10, 28, 60, 93, 5, 11, 40, 68, 25.

**Answer**


1. Insert 55 :




2. Insert 20 :




3. Insert 63 :




4. Insert 10 :




5. Insert 28 :




6. Insert 60 :




7. Insert 93 :




8. Insert 5 :




9. Insert 11 :




10. Insert 40 :



11. Insert 68 :



12. Insert 25 :



**PART-4**

*Graphs : Definition and Terminology, Representation of Graph.*

Questions-Answers

Long Answer Type and Medium Answer Type Questions

**Que 5.10.** What do you mean by graph? Also, explain directed and undirected graph.


**Answer**

A graph is a non-linear data structure consisting of nodes and edges. A graph consists of two sets as follows :

1. Set  $V$  of nodes or point or vertices of graph  $G$ .
2. Set  $E$  of ordered or unordered pairs of distinct edges of  $G$ .

We denote such a graph by  $G(V, E)$  and set of vertices as  $V(G)$  and set of edges as  $E(G)$ .

**For example :**




**Fig. 5.10.1.**

**Order :** If  $G$  is finite then number of vertices in  $G$  denoted by  $|V(G)|$  is called order of  $G$ .


**Size :** The number of edges denoted by  $|E(G)|$  in a finite graph  $G$  is called size of  $G$ .

**Directed graph :** A graph  $G(V, E)$  is said to be directed graph or digraph if each edge  $e \in E$  is associated with an ordered pair of vertices as shown below :



**Fig. 5.10.2.**

**Undirected graph :** A graph  $G(V, E)$  is said to be undirected if each edge  $e \in E$  is associated with an unordered pair of vertices as shown below :



**Fig. 5.10.3.**

**Que 5.11.** Discuss representation of graph.

**Answer**

Graph can be represented in following two ways :

**1. Matrix representation :**

Matrices are commonly used to represent graphs for computer processing. Advantages of representing the graph in matrix lies in the fact that many results of matrix algebra can be readily applied to study the structural properties of graph from an algebraic point of view.

**a. Adjacency matrix :**

**i. Representation of undirected graph :**

The adjacency matrix of a graph  $G$  with  $n$  vertices and no parallel edges is a  $n \times n$  matrix  $A = [a_{ij}]$  whose elements are given by


$$a_{ij} = 1, \text{ if there is an edge between } i^{\text{th}} \text{ and } j^{\text{th}} \text{ vertices} \\ = 0, \text{ if there is no edge between them}$$

**ii. Representation of directed graph :**

The adjacency matrix of a digraph  $D$ , with  $n$  vertices is the matrix

$$A = [a_{ij}]_{n \times n} \text{ in which} \\ a_{ij} = 1 \text{ if arc } (v_i, v_j) \text{ is in } D \\ = 0 \text{ otherwise}$$

**For example :**



**Fig. 5.11.1.**

$$A = \begin{matrix} & \begin{matrix} v_1 & v_2 & v_3 & v_4 \end{matrix} \\ \begin{matrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{bmatrix} \end{matrix}$$

**b. Incidence matrix :**

**i. Representation of undirected graph :**

Consider an undirected graph  $G = (V, E)$  which has  $n$  vertices and  $m$  edges all labelled. The incidence matrix  $I(G) = [b_{ij}]$  is then  $n \times m$  matrix, where


$$b_{ij} = 1 \text{ when edge } e_j \text{ is incident with } v_i \\ = 0 \text{ otherwise}$$

ii. **Representation of directed graph :**

The incidence matrix  $I(D) = [b_{ij}]$  of digraph  $D$  with  $n$  vertices and  $m$  edges is the  $n \times m$  matrix in which.

$$\begin{aligned} b_{ij} &= 1 \text{ if arc } j \text{ is directed away from a vertex } v_i \\ &= -1 \text{ if arc } j \text{ is directed towards vertex } v_i \\ &= 0 \text{ otherwise.} \end{aligned}$$

Find the incidence matrix to represent the graph shown in Fig. 5.11.2.



The incidence matrix of the digraph of Fig. 5.11.2 is

$$I(D) = \begin{bmatrix} 1 & 0 & 0 & -1 & 1 \\ -1 & 1 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & -1 \\ 0 & 0 & -1 & 1 & 0 \end{bmatrix}$$

2. **Linked representation :** In this representation, a list of vertices adjacent to each vertex is maintained. This representation is also called adjacency structure representation. In case of a directed graph, a care has to be taken, according to the direction of an edge, while placing a vertex in the adjacent structure representation of another vertex.

**PART-5**

*Multigraph, Bipartite Graphs, Planar Graph, Isomorphism and Homomorphism of Graphs.*

**Questions-Answers**

**Long Answer Type and Medium Answer Type Questions**

**Que 5.12.** Write short notes on :

- Simple and multigraph
- Complete graph and regular graph



- c. Bipartite graph
- d. Planar graph

**Answer**

**a. Simple and multigraph :**

i. **Simple graph :** A graph in which there is only one edge between a pair of vertices is called a simple graph.




Fig. 5.12.1.

ii. **Multigraph :** Any graph which contains some parallel edges is called a multigraph.




Fig. 5.12.2.

**b. Complete graph and regular graph :**

i. **Complete graph :** A simple graph, in which there is exactly one edge between each pair of distinct vertices is called a complete graph. The complete graph of  $n$  vertices is denoted by  $K_n$ . The graphs  $K_1$  to  $K_5$  are shown below in Fig. 5.12.3.




Fig. 5.12.3.

$K_n$  has exactly  $\frac{n(n-1)}{2} = {}^nC_2$  edges

ii. **Regular graph ( $n$ -regular graph) :** If every vertex of a simple graph has equal edges then it is called regular graph. If the degree of each vertex is  $n$  then the graph is called  $n$ -regular graph.




Fig. 5.12.4.

The graphs shown in Fig. 5.12.4 are 2-regular graphs.

The graph shown in Fig. 5.12.5 is 3-regular graph.



Fig. 5.12.5.

**Bipartite graph :**

- i. **Bipartite graph :** A graph  $G = (V, E)$  is bipartite if the vertex set  $V$  can be partitioned into two subsets (disjoint)  $V_1$  and  $V_2$  such that every edge in  $E$  connects a vertex in  $V_1$  and a vertex  $V_2$  (so that no edge in  $G$  connects either two vertices in  $V_1$  or two vertices in  $V_2$ ).  $(V_1, V_2)$  is called a bipartition of  $G$ .




Fig. 5.12.6. Some bipartite graphs.

- ii. **Complete bipartite graph :** The complete bipartite graph on  $m$  and  $n$  vertices, denoted  $K_{m, n}$  is the graph, whose vertex set is partitioned into sets  $V_1$  with  $m$  vertices and  $V_2$  with  $n$  vertices in which there is an edge between each pair of vertices  $v_1$  and  $v_2$  where  $v_1$  is in  $V_1$  and  $v_2$  is in  $V_2$ . The complete bipartite graphs  $K_{2,3}$ ,  $K_{2,4}$ ,  $K_{3,3}$ ,  $K_{3,5}$ , and  $K_{2,6}$





Fig. 5.12.7. Some complete bipartite graphs.

**Planar graph :**

A graph  $G$  is said to be planar if there exists some geometric representation of  $G$  which can be drawn on a plane such that no two of its edges intersect except only at the common vertex.

- i. A graph is said a planar graph, if it cannot be drawn on a plane without a crossover between its edges crossing.
- ii. The graphs shown in Fig. 5.12.8(a) and (b) are planar graphs.



(a)



(b)

Fig. 5.12.8. Some planar graph.

**Que 5.13.** Define the following with one example :

- i. Bipartite graph
- ii. Complete graph
- iii. How many edges in  $K_7$  and  $K_{3,6}$
- iv. Planar graph

AKTU 2017-18, Marks 10

**Answer**

- i. Refer Q. 5.12(c), Page 5-15C, Unit-5.
- ii. Refer Q. 5.12(b), Page 5-15C, Unit-5.
- iii. Number of edge in  $K_7$  : Since,  $K_n$  is complete graph with  $n$  vertices.

$$\text{Number of edge in } K_7 = \frac{7(7-1)}{2} = \frac{7 \times 6}{2} = 21$$

Number of edge in  $K_{3,6}$  :

Since,  $K_{n,m}$  is a complete bipartite graph with  $n \in V_1$  and  $m \in V_2$

$$\text{Number of edge in } K_{3,6} = 3 \times 6 = 18$$

- iv. Refer Q. 5.15(d), Page 5-18A, Unit-5.

**Que 5.14.** Explain isomorphism and homomorphism of graph.

**Answer**

**Isomorphism of graph :** Two graphs are isomorphic to each other if :

- i. Both have same number of vertices and edges.
- ii. Degree sequence of both graphs are same (degree sequence is the sequence of degrees of the vertices of a graph arranged in non-increasing order).

**Example :**



Fig. 5.14.1.

**Homomorphism of graph :** Two graphs are said to be homomorphic if one graph can be obtained from the other by the creation of edges in series (i.e., by insertion of vertices of degree two) or by the merger of edges in series.




Fig. 5.14.2.

**Que 5.15.** Prove that  $K_3$  and  $K_4$  are planar graphs. Prove that  $K_5$  is non-planar.

**Answer**

The complete  $K_3$  graph has 3 edges and 3 vertices.

For a graph to be planar  $3v - e \geq 6$

$$3v - e = 3 \times 3 - 3 = 9 - 3 = 6 \geq 6$$

$\therefore K_3$  is planar graph

Similarly complete  $K_4$  graph has 4 vertices and 6 edges.

$$3v - e = 3 \times 4 - 6 = 12 - 6 = 6 \geq 6$$

$\therefore K_4$  is planar graph

The complete  $K_5$  graph contains 5 vertices and 10 edges.

$$\text{Now } 3v - e = 3 \times 5 - 10 = 15 - 10 = 5 < 6$$

Hence  $K_5$  is non planar since for a graph to be planar  $3v - e \geq 6$ .

**Que 5.16.** What are Euler and Hamiltonian graph ?

OR

Explain the following terms with example :

- i. Homomorphism and Isomorphism graph
- ii. Euler graph and Hamiltonian graph
- iii. Planar and Complete bipartite graph

AKTU 2014-15, Marks 10

**Answer**

i. Refer Q. 5.14, Page 5-18C, Unit-5.

ii. **Eulerian path** : A path of graph  $G$  which includes each edge of  $G$  exactly once is called Eulerian path.

**Eulerian circuit** : A circuit of graph  $G$  which include each edge of  $G$  exactly once.

**Eulerain graph** : A graph containing an Eulerian circuit is called Eulerian graph.

**For example** : Graphs given below are Eulerian graphs.




Fig. 5.16.1.

**Hamiltonian graph :** A Hamiltonian circuit in a graph  $G$  is a closed path that visit every vertex in  $G$  exactly once except the end vertices. A graph  $G$  is called Hamiltonian graph if it contains a Hamiltonian circuit.

**For example :** Consider graphs given below :




Fig. 5.16.2.

Graph given is Fig. 5.16.2(a) is a Hamiltonian graph since it contains a Hamiltonian circuit  $A - B - C - D - A$  while graph in Fig 5.15.2(b) is not a Hamiltonian graph.

**Hamiltonian path :** The path obtained by removing any one edge from a Hamiltonian circuit is called Hamiltonian path. Hamiltonian path is subgraph of Hamiltonian circuit. But converse is not true.

The length of Hamiltonian path in a connected graph of  $n$  vertices is  $n - 1$  if it exists.

iii. Refer Q. 5.12, Page 5-15C, Unit-5.

**Que 5.17.**

- Prove that a connected graph  $G$  is Euler graph if and only if every vertex of  $G$  is of even degree.
- Which of the following simple graph have a Hamiltonian circuit or, if not a Hamiltonian path ?

AKTU 2016-17, Marks 15




Fig. 5.17.1.

**Answer**

- First of all we shall prove that if a non-empty connected graph is Eulerian then it has no vertices of odd degree.
  - Let  $G$  be Eulerian.
  - Then  $G$  has an Eulerian trail which begins and ends at  $u$ .

4. If we travel along the trail then each time we visit a vertex. We use two edges, one in and one out.
5. This is also true for the start vertex because we also end there.
6. Since an Eulerian trail uses every edge once, the degree of each vertex must be a multiple of two and hence there are no vertices of odd degree.
7. Now we shall prove that if a non-empty connected graph has no vertices of odd degree then it is Eulerian.
8. Let every vertex of  $G$  have even degree.
9. We will now use a proof by mathematical induction on  $|E(G)|$ , the number of edges of  $G$ .

**Basis of induction :**

Let  $|E(G)| = 0$ , then  $G$  is the graph  $K_1$ , and  $G$  is Eulerian.

**Inductive step :**

1. Let  $P(n)$  be the statement that all connected graphs on  $n$  edges of even degree are Eulerian.
2. Assume  $P(n)$  is true for all  $n < |E(G)|$ .
3. Since each vertex has degree at least two,  $G$  contains a cycle  $C$ .
4. Delete the edges of the cycle  $C$  from  $G$ .
5. The resulting graph,  $G'$  say, may not be connected.
6. However, each of its components will be connected, and will have fewer than  $|E(G)|$  edges.
7. Also, all vertices of each component will be of even degree, because the removal of the cycle either leaves the degree of a vertex unchanged, or reduces it by two.
8. By the induction assumption, each component of  $G'$  is therefore Eulerian.
9. To show that  $G$  has an Eulerian trail, we start the trail at a vertex,  $u$  say, of the cycle  $C$  and traverse the cycle until we meet a vertex,  $c_1$  say, of one of the components of  $G'$ .
10. We then traverse that component's Eulerian trail, finally returning to the cycle  $C$  at the same vertex,  $c_1$ .
11. We then continue along the cycle  $C$ , traversing each component of  $G'$  as it meets the cycle.
12. Eventually, this process traverses all the edges of  $G$  and arrives back at  $u$ , thus producing an Eulerian trail for  $G$ .
13. Thus,  $G$  is Eulerian by the principle of mathematical induction.

**G1 :** The graph  $G_1$  shown in Fig. 5.17.1 contains Hamiltonian circuit, i.e.,  $a - b - c - d - e - a$  and also a Hamiltonian path, i.e.,  $abcde$ .

**G2 :** The graph  $G_2$  shown in Fig. 5.17.1 does not contain Hamiltonian circuit since every cycle containing every vertex must contain the edge  $e$  twice. But the graph does have a Hamiltonian path  $a - b - c - d$ .

**G3:** The graph G3 shown in Fig. 5.17.1 neither have Hamiltonian circuit nor have Hamiltonian path because any traversal does not cover all the vertices.

**Que 5.18.** Prove that a simple graph with  $n$  vertices and  $k$  components can have at most  $\frac{(n-k)(n-k+1)}{2}$  edges.

AKTU 2016-17, Marks 10

**Answer**

Let the number of vertices in each of the  $k$ -components of a graph  $G$  be  $n_1, n_2, \dots, n_k$ , then we get

$$n_1 + n_2 + \dots + n_k = n \text{ where } n_i \geq 1 (i = 1, 2, \dots, k)$$

Now, 
$$\sum_{i=1}^k (n_i - 1) = \sum_{i=1}^k n_i - \sum_{i=1}^k 1 = n - k$$

$$\therefore \left( \sum_{i=1}^k (n_i - 1) \right)^2 = n^2 + k^2 - 2nk$$

$$\text{or } \sum_{i=1}^k (n_i - 1)^2 + 2 \sum_{\substack{i=1 \\ i \neq j}}^k \sum_{j=1}^k (n_i - 1)(n_j - 1) = n^2 + k^2 - 2nk$$

$$\text{or } \sum_{i=1}^k (n_i - 1)^2 + 2(\text{non-negative terms}) = n^2 + k^2 - 2nk$$

$$[\because n_i - 1 \geq 0, n_j - 1 \geq 0]$$

$$\text{or } \sum_{i=1}^k (n_i - 1)^2 \leq n^2 + k^2 - 2nk$$

$$\text{or } \sum_{i=1}^k n_i^2 + \sum_{i=1}^k 1 - 2 \sum_{i=1}^k n_i \leq n^2 + k^2 - 2nk$$

$$\text{or } \sum_{i=1}^k n_i^2 + k - 2n \leq n^2 + k^2 - 2nk$$

$$\text{or } \sum_{i=1}^k n_i^2 - n \leq n^2 + k^2 - 2nk - k + n$$

$$= n(n - k + 1) - k(n - k + 1)$$

$$= (n - k)(n - k + 1) \dots (5.18.1)$$

We know that the maximum number of edges in the  $i^{\text{th}}$  component of

$$G = {}^n C_2 = \frac{n_i(n_i - 1)}{2}$$

Therefore, the maximum number of edges in  $G$  is :

$$\frac{1}{2} \sum n_i(n_i - 1) = \frac{1}{2} (\sum n_i^2 - \sum n_i) = \frac{1}{2} (\sum n_i^2 - n)$$

$$\leq \frac{1}{2} (n - k)(n - k + 1) \text{ by using eq. (5.18.1)}$$

**Que 5.19.** What are different ways to represent a graph? Define Euler circuit and Euler graph. Give necessary and sufficient conditions for Euler circuits and paths. **AKTU 2018-19, Marks 07**

**Answer**

**Representation of graph :** Refer Q. 5.11, Page 5-14C, Unit-5.

**Euler circuit and Euler graph :** Refer Q. 5.16, Page 5-19C, Unit-5.

**Necessary and sufficient condition for Euler circuits and paths :**

1. A graph has an Euler circuit if and only if the degree of every vertex is even.
2. A graph has an Euler path if and only if there are at most two vertices with odd degree.

**Que 5.20.** Define and explain any two the following :

1. BFS and DFS in trees
2. Euler graph
3. Adjacency matrix of a graph

**AKTU 2017-18, Marks 07**

**Answer**

1. **Breadth First Search (BFS) :** Breadth First Search (BFS) is an algorithm for traversing or searching tree or graph data structures. It starts at the tree root and explores the neighbour nodes first, before moving to the next level neighbours.

**Algorithmic steps :**

**Step 1 :** Push the root node in the queue.

**Step 2 :** Loop until the queue is empty.

**Step 3 :** Remove the node from the queue.

**Step 4 :** If the removed node has unvisited child nodes, mark them as visited and insert the unvisited children in the queue.

**Depth First Search (DFS) :**

Depth First Search (DFS) is an algorithm for traversing or searching tree or graph data structures. One starts at the root (selecting some arbitrary node as the root in the case of a graph) and explores as far as possible along each branch before backtracking.

**Algorithmic steps :**

**Step 1 :** Push the root node in the stack.

**Step 2 :** Loop until stack is empty.

**Step 3 :** Pick the node of the stack.

**Step 4 :** If the node has unvisited child nodes, get the unvisited child node, mark it as traversed and push it on stack.

Step 5 : If the node does not have any unvisited child nodes, pop the node from the stack.

2. Refer Q. 5.16, Page 5-19C, Unit-5.
3. Refer Q. 5.11, Page 5-14C, Unit-5.

## PART-6

### Graph Coloring.

#### Questions-Answers

#### Long Answer Type and Medium Answer Type Questions

**Que 5.21.** Write a short note on graph coloring.

#### Answer

1. Suppose that  $G(V, E)$  is a graph with no multiple edges, a vertex colouring of  $G$  is an assignment of colours.
2. A graph  $G$  is  $m$ -colourable if there exists a colouring of  $G$  which uses  $m$  colours.
3. Colouring the vertices such a way such that no two adjacent vertices have same colour is called proper colouring otherwise it is called improper colouring.

## PART-7

*Recurrence Relation and Generating Function : Recursive Definition of Function, Recursive Algorithms, Method of Solving Recurrences, Combinatorics : Introduction, Counting Techniques, Pigeonhole Principle.*

#### Questions-Answers

#### Long Answer Type and Medium Answer Type Questions

**Que 5.22.** Write a short note on recurrence relation.

#### Answer

For a sequence of numbers or numeric function  $(a_0, a_1, a_2, \dots, a_r, \dots)$  an equation relating  $a_r$ , for any  $r$ , to one or more of the  $a_i$ 's,  $i < r$  is called

recurrence relation. Recurrence relations are also called difference equations because they can be written in terms of the difference between the consecutive terms of a sequence.

**For example :**

$$\begin{aligned} a_r &= 4a_{r-1} & r \geq 1 \\ a_r &= a_{r-1} - 2a_{r-2} & r \geq 2 \end{aligned}$$

are recurrence relations.

**Order of recurrence relation :** The difference between the highest and lowest subscripts of  $a_r$  or  $f(x)$  or  $y_n$  is called the order of recurrence relation.

**For example :**

1. The equation  $13a_r + 12a_{r-1} = 0$  is of order 1 (i.e.,  $r - (r - 1) = 1$ ).
2. The equation  $8f(x) + 2f(x + 1) + f(x + 2) = k(x)$  is of order 2 (i.e.,  $(x + 2) - x = 2$ ).

**Degree of recurrence relation :** The highest power of  $a_r$  or  $f(x)$  or  $y_n$  is called degree of recurrence relation.

**For example :**

1. The equation  $y_{k+2}^3 + 2y_{k+1}^2 - y_k = 0$  has the degree 3 as the highest power of  $y_k$  is 3.
2. The equation  $a_r^4 + 2a_{r-1}^3 + 3a_{r-2}^2 + 4a_{r-3} = 0$  has the degree 4, as the highest power of  $a_r$  is 4.

**Que 5.23.** What do you mean generating function ? Solve the

recurrence relation :

$$a_n = 2a_{n-1} - a_{n-2}, n \geq 2 \text{ given } a_0 = 3, a_1 = -2$$

using generating function.

**Answer**

**Generating function :** The generating function for the sequence  $a_0, a_1, \dots, a_k, \dots$  of real numbers is infinite series given by

$$G(x) = a_0 + a_1x + a_2x^2 + \dots + a_kx^k + \dots = \sum_{k=0}^{\infty} a_kx^k$$

$x$  is considered just a symbol called indeterminate and it is not variable, which is replaced by numbers belonging to same domain.

The given recurrence relation is,

$$a_n = 2a_{n-1} - a_{n-2}, n \geq 2 \quad \dots(5.23.1)$$

Multiply by  $x^n$  and take summation from  $n = 2$  to  $\infty$ , we get

$$\sum_{n=2}^{\infty} a_n x^n = 2 \sum_{n=2}^{\infty} a_{n-1} x^n - \sum_{n=2}^{\infty} a_{n-2} x^n \quad \dots(5.23.2)$$

We know,

$$G(x) = \sum_{n=0}^{\infty} a_n x^n = a_0 + a_1x + a_2x^2 + a_3x^3 + \dots \quad \dots(5.23.3)$$

From eq. (5.23.2), we have

$$(a_2x^2 + a_3x^3 + \dots) = 2(a_1x^2 + a_2x^3 + \dots) - (a_0x^2 + a_1x^3 + \dots)$$

$$(a_2x^2 + a_3x^3 + \dots) = 2x(a_1x + a_2x^2 + \dots) - x^2(a_0 + a_1x + \dots)$$

Using eq. (5.23.3), we get

$$G(x) - a_0 - a_1x = 2x(G(x) - a_0) - x^2 G(x)$$

$$G(x) - 3 + 2x = 2x(G(x) - 3) - x^2 G(x)$$

$$G(x)[1 - 2x + x^2] = 3 - 8x$$

$$G(x) = \frac{3 - 8x}{x^2 - 2x + 1} = \frac{3 - 8x}{(x - 1)^2} = \frac{3 - 8x}{(1 - x)^2} = \frac{3}{(1 - x)^2} - \frac{8x}{(1 - x)^2}$$

$$\therefore a_n = 3(n + 1) - 8n = 3 - 5n$$

**Que 5.24.** Solve the recurrence relation  $y_{n+2} - 5y_{n+1} + 6y_n = 5^n$

subject to the condition  $y_0 = 0, y_1 = 2$ .

**AKTU 2016-17, Marks 10**

**Answer**

Let  $G(t) = \sum_{n=0}^{\infty} a_n t^n$  be generating function of sequence  $\{a_n\}$ .

Multiplying given equation by  $t^n$  and summing from  $n = 0$  to  $\infty$ , we have

$$\sum_{n=0}^{\infty} a_{n+2} t^n - 5 \sum_{n=0}^{\infty} a_{n+1} t^n + 6 \sum_{n=0}^{\infty} a_n t^n = \sum_{n=0}^{\infty} 5^n t^n$$

$$\frac{G(t) - a_0 - a_1 t}{t^2} - 5 \left[ \frac{G(t) - a_0}{t} \right] + 6G(t) = \frac{1}{1 - 5t}$$

Put  $a_0 = 0$  and  $a_1 = 2$

$$G(t) - 2t - 5t G(t) + 6t^2 G(t) = \frac{t^2}{1 - 5t}$$

$$G(t) - 5t G(t) + 6t^2 G(t) = \frac{t^2}{1 - 5t} + 2t$$

$$G(t) (1 - 5t + 6t^2) = \frac{t^2}{1 - 5t} + 2t$$

$$(6t^2 - 5t + 1) G(t) = \frac{t^2}{1 - 5t} + 2t$$

$$G(t) = \frac{t^2}{(1 - 5t)(3t - 1)(2t - 1)} + \frac{2t}{(3t - 1)(2t - 1)}$$

$$= \frac{t^2}{(1 - 5t)(1 - 3t)(1 - 2t)} + \frac{2t}{(1 - 3t)(1 - 2t)}$$

Let

$$\frac{t^2}{(1 - 5t)(1 - 3t)(1 - 2t)} = \frac{A}{(1 - 5t)} + \frac{B}{(1 - 3t)} + \frac{C}{(1 - 2t)}$$

$$A = (1-5t) \frac{t^2}{(1-5t)(1-3t)(1-2t)} \Big|_{t=1/5}$$

$$= \frac{t^2}{(1-3t)(1-2t)} \Big|_{t=1/5}$$

$$= \frac{1/25}{(1-3/5)(1-2/5)} = \frac{1}{6}$$

$$B = (1-3t) \frac{t^2}{(1-5t)(1-3t)(1-2t)} \Big|_{t=1/3}$$

$$= \frac{t^2}{(1-5t)(1-2t)} \Big|_{t=1/3} = \frac{1/9}{\left(\frac{3-5}{3}\right)\left(\frac{3-2}{3}\right)}$$

$$= -\frac{1}{2}$$

$$C = (1-2t) \frac{t^2}{(1-5t)(1-3t)(1-2t)} \Big|_{t=1/2}$$

$$= \frac{t^2}{(1-5t)(1-3t)} \Big|_{t=1/2} = \frac{1/4}{\frac{(2-5)}{2} \times \frac{(2-3)}{2}}$$

$$= \frac{1}{3}$$

Again,

$$\frac{2t}{(1-3t)(1-2t)} = \frac{D}{(1-3t)} + \frac{E}{(1-2t)}$$

$$D = (1-3t) \frac{2t}{(1-3t)(1-2t)} \Big|_{t=1/3}$$

$$= \frac{2t}{(1-2t)} \Big|_{t=1/3} = \frac{2/3}{(3-2)} = 2$$

$$E = (1-2t) \frac{2t}{(1-3t)(1-2t)} \Big|_{t=1/2}$$

$$= \frac{2t}{(1-3t)} \Big|_{t=1/2} = \frac{2/2}{\frac{2-3}{2}} = -2$$

$$G(t) = \frac{1/6}{(1-5t)} - \frac{1/2}{(1-3t)} + \frac{1/3}{(1-2t)} + \frac{2}{(1-3t)} - \frac{2}{(1-2t)}$$

$$= \frac{1/6}{1-5t} + \frac{3/2}{(1-3t)} - \frac{5/3}{1-2t}$$

$$\sum_{n=0}^{\infty} a_n t^n = \frac{1}{6} \sum_{n=0}^{\infty} (5t)^n + \frac{3}{2} \sum_{n=0}^{\infty} (3t)^n - \frac{5}{3} \sum_{n=0}^{\infty} (2t)^n$$

$$\therefore a_n = \frac{1}{6}(5)^n + \frac{3}{2}(3)^n - \frac{5}{3}(2)^n$$

**Que 5.25.** Solve the recurrence relation by the method of generating function :

$$a_r - 7a_{r-1} + 10a_{r-2} = 0, r \geq 2. \text{ Given } a_0 = 3 \text{ and } a_1 = 3.$$

AKTU 2014-15, Marks 10

OR

Solve the recurrence relation using generating function :

$$a_n - 7a_{n-1} + 10a_{n-2} = 0 \text{ with } a_0 = 3, a_1 = 3.$$

AKTU 2015-16, Marks 10

Answer

$$a_r - 7a_{r-1} + 10a_{r-2} = 0, r \geq 2$$

Multiply by  $x^r$  and take sum from 2 to  $\infty$ .

$$\sum_{r=2}^{\infty} a_r x^r - 7 \sum_{r=2}^{\infty} a_{r-1} x^r + 10 \sum_{r=2}^{\infty} a_{r-2} x^r = 0$$

$$(a_2 x^2 + a_3 x^3 + a_4 x^4 + \dots) - 7(a_1 x^2 + a_2 x^3 + \dots)$$

$$+ 10(a_0 x^2 + a_1 x^3 + \dots) = 0$$

We know that

$$G(x) = \sum_{r=0}^{\infty} a_r x^r = a_0 + a_1 x + \dots$$

$$G(x) - a_0 - a_1 x - 7x(G(x) - a_0) + 10x^2 G(x) = 0$$

$$G(x) [1 - 7x + 10x^2] = a_0 + a_1 x - 7a_0 x$$

$$= 3 + 3x - 21x = 3 - 18x$$

$$G(x) = \frac{3 - 18x}{10x^2 - 7x + 1} = \frac{3 - 18x}{10x^2 - 5x - 2x + 1}$$

$$= \frac{3 - 18x}{5x(2x - 1) - 1(2x - 1)} = \frac{3 - 18x}{(5x - 1)(2x - 1)}$$

Now,

$$\frac{3 - 18x}{(5x - 1)(2x - 1)} = \frac{A}{5x - 1} + \frac{B}{2x - 1}$$

$$3 - 18x = A(2x - 1) + B(5x - 1)$$

put  $x = \frac{1}{2}$

$$3 - 9 = B \left( \frac{5}{2} - 1 \right) \Rightarrow -6 = \frac{3}{2} B \Rightarrow B = -4$$

put  $x = \frac{1}{5}$

$$3 - \frac{18}{5} = A \left( \frac{2}{5} - 1 \right) \Rightarrow -\frac{3}{5} = -\frac{3}{5} A = 1 \Rightarrow A = 1$$

$$\therefore G(x) = \frac{1}{5x-1} - \frac{4}{2x-1} = \frac{4}{1-2x} - \frac{1}{1-5x}$$

$$\therefore a^r = 4 \cdot 2^r - 5^r$$

**Que 5.26.** Solve the recurrence relation  $a_{r+2} - 5a_{r+1} + 6a_r = (r+1)^2$ .

**AKTU 2014-15, Marks 10**

**Answer**

$$a_{r+2} - 5a_{r+1} + 6a_r = (r+1)^2 = r^2 + 2r + 1 \quad \dots(5.26.1)$$

Now the characteristic equation is :

$$x^2 - 5x + 6 = 0$$

$$(x-3)(x-2) = 0 \Rightarrow x = 3, 2$$

The homogeneous solution is :

$$a_r^{(h)} = C_1 2^r + C_2 3^r$$

Let the particular solution be :

$$a_r^{(p)} = A_0 + A_1 r + A_2 r^2$$

From eq. (5.26.1)

$$\begin{aligned} A_0 + A_1(r+2) + A_2(r+2)^2 - 5[A_0 + A_1(r+1)] + A_2(r+1)^2 \\ = r^2 + 2r + 1 \end{aligned}$$

$$\begin{aligned} (A_0 + 2A_1 + 4A_2 - 5A_0 - 5A_1 - 5A_2 + 6A_0) + r(A_1 + 4A_2 - 5A_1 - 10A_2 + 6A_1) \\ + r^2(A_2 - 5A_2 + 6A_2) = r^2 + 2r + 1 \end{aligned}$$

Comparing both sides, we get,

$$2A_0 - 3A_1 - A_2 = 1 \quad \dots(5.26.2)$$

$$2A_1 - 6A_2 = 2 \quad \dots(5.26.3)$$

$$2A_2 = 1 \Rightarrow A_2 = 1/2$$

From eq. (5.26.3),  $2A_1 - 3 = 2$

$$A_1 = \frac{5}{2}$$

From eq. (5.26.2)

$$2A_0 - \frac{15}{2} - \frac{1}{2} = 1$$

$$2A_0 - 8 = 1 \Rightarrow A_0 = \frac{9}{2}$$

$$a_r^{(p)} = \frac{9}{2} + \frac{5}{2}r + \frac{r^2}{2}$$

The final solution is,  $a_r = a_r^{(h)} + a_r^{(p)} = C_1 2^r + C_2 3^r + \frac{9}{2} + \frac{5}{2}r + \frac{r^2}{2}$

**Que 5.27.** Solve  $a_r - 6a_{r-1} + 8a_{r-2} = r \cdot 4^r$ , given  $a_0 = 8$ , and  $a_1 = 1$ .

**AKTU 2017-18, Marks 07**

**Answer**

$$a_r - 6a_{r-1} + 8a_{r-2} = r \cdot 4^r$$

The characteristic equation is,  $x^2 - 6x + 8 = 0$ ,  $x^2 - 2x - 4x + 8 = 0$   
 $(x - 2)(x - 4) = 0$ ,  $x = 2, 4$

The solution of the associated non-homogeneous recurrence relation is,

$$a_r^{(h)} = B_1(2)^r + B_2(4)^r \quad \dots(5.27.1)$$

Let particular solution of given equation is,  $a_r^{(p)} = r^2(A_0 + A_1r)4^r$

Substituting in the given equation, we get

$$\Rightarrow r^2(A_0 + A_1r)4^r - 6(r-1)^2(A_0 + A_1(r-1))4^{r-1} + 8(r-2)^2(A_0 + A_1(r-2))4^{r-2} = r \cdot 4^r$$

$$\Rightarrow r^2A_0 + A_1r^3 - \frac{6}{4} [(A_0r^2 - 2A_0r + A_0) + (A_1r^3 - A_1 - 3A_1r^2 + 3A_1r)^2] + \frac{8}{4^2} [(A_0r^2 - 4rA_0 + 4A_0) + (A_1r^3 - 8A_1 - 6A_1r^2 + 12A_1r)] = r$$

$$\Rightarrow rA_0 + A_1r^3 - \frac{3}{2}A_0r^2 + 3A_0r - \frac{3}{2}A_0 - \frac{3}{2}A_1r^3 + \frac{3}{2}A_1 + \frac{9}{2}A_1r^2 - \frac{9}{2}A_1r + \frac{1}{2}A_0r^2 - 2A_0r + 2A_0 + \frac{1}{2}A_1r^3 - 4A_1 - 3A_1r^2 - 6A_1r = r$$

$$\Rightarrow 2A_0r - A_0r^2 - \frac{1}{2}A_0 - \frac{5}{2}A_1 + \frac{3}{2}A_1r^2 + \frac{3}{2}A_1r = r$$

Comparing both sides, we get

$$2A_0 + \frac{3}{2}A_1 = 1 \quad \dots(5.27.2)$$

$$A_0 + 5A_1 = 0 \quad \dots(5.27.3)$$

Solving equation (5.27.2) and (5.27.3), we get  $A_1 = \frac{-2}{17}$   $A_0 = \frac{-10}{17}$

To find the value of  $B_1$  and  $B_2$  put  $r = 0$  and  $r = 1$  in equation (5.27.1)

$$r = 0 \quad a_0 = B_1 + B_2 \quad B_1 + B_2 = 8 \quad \dots(5.27.4)$$

$$r = 1 \quad a_1 = 2B_1 + 4B_2 \quad 2B_1 + 4B_2 = 1 \quad \dots(5.27.5)$$

Solving equations (5.27.4) and (5.27.5), we get  $B_1 = \frac{31}{2}$   $B_2 = \frac{-15}{2}$

Complete solution is,  $a_r = a_r^{(h)} + a_r^{(p)}$

$$a_r = \frac{31}{2} 2^r - \frac{15}{2} 4^r + r^2 \left[ \left( \frac{-10}{17} \right) + \left( \frac{-2}{17} \right) r \right] 4^r$$

**Que 5.28.** Solve the recurrence relation :  $a_r + 4a_{r-1} + 4a_{r-2} = r^2$ .

**AKTU 2017-18, Marks 07**

**Answer**

$$a_r + 4a_{r-1} + 4a_{r-2} = r^2$$

The characteristic equation is,

$$x^2 + 4x + 4 = 0$$

$$(x + 2)^2 = 0$$

$$x = -2, -2$$

The homogeneous solution is,  $a^{(h)} = (A_0 + A_1 r) (-2)^r$

The particular solution be,  $a^{(p)} = (A_0 + A_1 r) r^2$

Put  $a_r, a_{r-1}$  and  $a_{r-2}$  from  $a^{(p)}$  in the given equation, we get

$$r^2 A_0 + A_1 r^3 + 4A_0 (r-1)^2 + 4A_1 (r-1)^3 + 4A_0 (r-2)^2 + 4A_1 (r-2)^3 = r^2$$

$$A_0 (r^2 + 4r^2 - 8r + 4 + 4r^2 - 16r + 16) +$$

$$A_1 (r^3 + 4r^3 - 4 - 12r^2 + 12r + 4r^3 - 32 - 24r^2 + 48r) = r^2$$

$$A_0 (9r^2 - 24r + 20) + A_1 (9r^3 - 48r^2 + 60r - 36) = r^2$$

Comparing the coefficient of same power of  $r$ , we get

$$9A_0 - 48A_1 = 1 \quad \dots(5.28.1)$$

$$20A_0 - 36A_1 = 0 \quad \dots(5.28.2)$$

Solving equation (5.28.1) and (5.28.2)  $A_0 = \frac{-3}{53}$   $A_1 = \frac{-5}{159}$

The complete solution is,

$$a_r = a_r^{(p)} + a_r^{(h)} = (A_0 + A_1 r) (-2)^r + \left[ \left( \frac{-3}{53} \right) + \left( \frac{-5}{159} \right) r \right] r^2$$

**Que 5.29.** Suppose that a valid codeword is an  $n$ -digit number in decimal notation containing an even number of 0's. Let  $a_n$  denote the number of valid codewords of length  $n$  satisfying the recurrence relation  $a_n = 8a_{n-1} + 10^{n-1}$  and the initial condition  $a_1 = 9$ . Use generating functions to find an explicit formula for  $a_n$ .

**AKTU 2018-19, Marks 07**

**Answer**

Let  $G(x) = \sum_{n=0}^{\infty} a_n x^n$  be the generating function of the sequence  $a_0, a_1, a_2, \dots$

we sum both sides of the last equations starting with  $n = 1$ . To find that

$$\begin{aligned} G(x) - 1 &= \sum_{n=1}^{\infty} a_n x^n = \sum_{n=1}^{\infty} (8a_{n-1}x^n + 10^{n-1}x^n) \\ &= 8 \sum_{n=1}^{\infty} a_{n-1}x^n + \sum_{n=1}^{\infty} 10^{n-1}x^n \\ &= 8x \sum_{n=1}^{\infty} a_{n-1}x^{n-1} + x \sum_{n=1}^{\infty} 10^{n-1}x^{n-1} \\ &= 8x \sum_{n=0}^{\infty} a_n x^n + x \sum_{n=0}^{\infty} 10^n x^n \\ &= 8xG(x) + x/(1 - 10x) \end{aligned}$$

Therefore, we have

$$G(x) - 1 = 8xG(x) + x/(1 - 10x)$$

Expanding the right hand side of the equation into partial fractions gives

$$G(x) = \frac{1}{2} \left( \frac{1}{1 - 8x} + \frac{1}{1 - 10x} \right)$$

This is equivalent to

$$\begin{aligned} G(x) &= \frac{1}{2} \left( \sum_{n=0}^{\infty} 8^n x^n + \sum_{n=0}^{\infty} 10^n x^n \right) \\ &= \sum_{n=0}^{\infty} \frac{1}{2} (8^n + 10^n) x^n \\ a_n &= \frac{1}{2} (8^n + 10^n) \end{aligned}$$

**Que 5.30.** Define permutation and combination. Also, write difference between them.

**Answer**

1. Permutation refers to different ways of arranging a set of object in a sequential order.
2. The number of permutations of  $n$  different things taken  $r$  ( $\leq n$ ) at a time is denoted by  $p(n, r)$  or  ${}^n P_r$ .
3. Combination refers to several ways of choosing items from a large set of object.

4. The number of combinations of  $n$  different thing taken  $r$  ( $\leq n$ ) at a time is denoted by  $C(n, r)$  or  ${}^n C_r$

The selection of two letters from three letters  $a, b, c$  are  $ab, bc, ca$  and thus, the number of combinations of 3 letters taken 2 at a time is  $C(3, 2) = 3$

**Difference between a permutation and combination :**

S. No.	Permutation	Combination
1.	Both selection and arrangement are made.	Only selection is made.
2.	Ordering of the selected object is essential.	Ordering of the selected object is not essential.
3.	Multiple permutations can be derive from combination.	Single combination is derive from single permutation.
4.	${}^n P_r = \frac{n!}{(n-r)!}$	${}^n C_r = \frac{n!}{r!(n-r)!}$

**Que 5.31.** Suppose that a cookie shop has four different kinds of cookies. How many different ways can six cookies be chosen ?

**AKTU 2016-17, Marks 15**

**Answer**

As the order in which each cookie is chosen does not matter and each kind of cookies can be chosen as many as 6 times, the number of ways these cookies can be chosen is the number of 6-combination with repetition allowed from a set with 4 distinct elements.

The number of ways to choose six cookies in the bakery shop is the number of 6 combinations of a set with four elements.

$$C(4 + 6 - 1, 6) = C(9, 6)$$

Since  $C(9, 6) = C(9, 3) = (9 \cdot 8 \cdot 7) / (1 \cdot 2 \cdot 3) = 84$

Therefore, there are 84 different ways to choose the six cookies.

**Que 5.32.** Write short notes on the following :

- Recursive algorithms
- Pigeonhole principle

**Answer**

- Recursive algorithm :

A function is called recursively defined if the function refers to itself, and satisfies the following step :

- There must be certain arguments for which the function does not refer to itself, these arguments are called initial values (Base values).



# Exclusive training



5-34 C (CS/IT-Sem-3)

Trees, Graph and Combinatorics

b. These base values are used to define the other values of the function by using the function recursively.

**Example :**  $a_r = 3 a_{r-1}, r \geq 1, a_0 = 1$   
 then  $a_1 = 3a_0 = 3$   
 $a^2 = 3a_1 = 3(3a_0) = 3^2 a_0 = 3^2$

and so on.

$$a_r = 3^r, r \geq 0$$

ii. **Pigeonhole principle :**

The pigeonhole principle is sometime useful in counting methods.

If  $n$  pigeons are assigned to  $m$  pigeonholes then at least one pigeonhole contains two or more pigeons ( $m < n$ ).

**Proof :**

1. Let  $m$  pigeonholes be numbered with the numbers 1 through  $m$ .
2. Beginning with the pigeon 1, each pigeon is assigned in order to the pigeonholes with the same number.
3. Since  $m < n$  i.e., the number of pigeonhole is less than the number of pigeons,  $n-m$  pigeons are left without having assigned a pigeonhole.
4. Thus, at least one pigeonhole will be assigned to a more than one pigeon.
5. We note that the pigeonhole principle tells us nothing about how to locate the pigeonhole that contains two or more pigeons.
6. It only asserts the existence of a pigeon hole containing two or more pigeons.
7. To apply the principle one has to decide which objects will play the role of pigeon and which objects will play the role of pigeonholes.

**Que 5.33.** Find the number of integers between 1 and 250 that are divisible by any of the integers 2, 3, 5, and 7.

**Answer**

1, 2, 3, ..... 250

Number of integers between 1 and 250 that are divisible by 2 :

Quotient of last number  $\div 2$  - quotient of first number  $\div 2$

$$(250 \div 2) - (1 \div 2) = 125 - 0 = 125$$

Number of integers between 1 and 250 that are divisible by 3

$$(250 \div 3) - (1 \div 3) = 83 - 0 = 83$$

Number of integers between 1 and 250 that are divisible by 5

$$(250 \div 5) - (1 \div 5) = 50 - 0 = 50$$

Number of integers between 1 and 250 that are divisible by 7

$$(250 \div 7) - (1 \div 7) = 35 - 0 = 5$$

$\therefore$  Total number of integers divisible by only 2, 3, 5, 7, individually are

$$125 + 83 + 50 + 35 = 293$$

Number of integers divisible by (2 and 3) = 41

Downloaded from



Number of integers divisible by (2 and 5) = 25

Number of integers divisible by (2 and 7) = 17

Number of integers divisible by (3 and 5) = 16

Number of integers divisible by (3 and 7) = 11

Number of integers divisible by (5 and 7) = 7

Number of integers divisible by (2, 3 and 5) = 8

Number of integers divisible by (2, 3, and 7) = 5

Number of integers divisible by (2, 5, and 7) = 3

Number of integers divisible by (3, 5 and 7) = 2

135

∴ Number of integers between 1 and 250 that are divisible by any of the integer (2, 3, 5 and 7) =  $293 - 135 = 158$ .

**Que 5.34.** How many different rooms are needed to assign 500 classes, if there are 45 different time periods during in the university time table that are available ?

**Answer**

Using pigeonhole principle :

Here  $n = 500, m = 45 = \left\lceil \frac{n-1}{m} \right\rceil + 1 = \left\lceil \frac{500-1}{45} \right\rceil + 1$

At least 12 rooms are needed.

**Que 5.35.** A total of 1232 student have taken a course in Spanish, 879 have taken a course in French, and 114 have taken a course in Russian. Further 103 have taken courses in both Spanish and French, 23 have taken courses in both Spanish and Russian, and 14 have taken courses in both French and Russian. If 2092 students have taken least one of Spanish, French and Russian, how many students have taken a course in all three languages ?

**AKTU 2018-19, Marks 07**

**Answer**

Let  $S$  be the set of students who have taken a course in Spanish,  $F$  be the set of students who have taken a course in French, and  $R$  be the set of students who have taken a course in Russian. Then, we have

$$|S| = 1232, |F| = 879, |R| = 114, |S \cap F| = 103, |S \cap R| = 23, \\ |S \cap R| = 14, \text{ and } |S \cup F \cup R| = 2092.$$

Using the equation

$$|S \cup F \cup R| = |S| + |F| + |R| - |S \cap F| - |S \cap R| - |S \cap R| + |S \cap F \cap R|,$$

5-36 C (CS/IT-Sem-3)

Trees, Graph and Combinatorics

$$2092 = 1232 + 879 + 114 - 103 - 23 - 14 + |S \cap F \cap R|,$$

$$|S \cap F \cap R| = 7.$$




Fig. 5.35.1.

**VERY IMPORTANT QUESTIONS**

Following questions are very important. These questions may be asked in your **SESSIONALS** as well as **UNIVERSITY EXAMINATION**.

Q.1. Explain in detail about the binary tree traversal with an example.

ANS: Refer Q. 5.3.

Q.2. Define a binary tree. A binary tree has 11 nodes. It's inorder and preorder traversals node sequences are :

Preorder : **ABDHIEJL CFG**

Inorder : **HDIBJEKAF CG**

Draw the tree.

ANS: Refer Q. 5.5.

Q.3. Given the inorder and postorder traversal of a tree T:

Inorder : **HFEABIGDC** Postorder : **BEHFACDGI**

Determine the tree T and it's Preorder.

ANS: Refer Q. 5.6.

Q. 4. Define the following with one example :

- i. Bipartite graph
- ii. Complete graph
- iii. How many edges in  $K_7$  and  $K_{3,0}$
- iv. Planar graph

**Ans.** Refer Q. 5.13.


Q. 5. Explain the following terms with example :

- i. Homomorphism and Isomorphism graph
- ii. Euler graph and Hamiltonian graph
- iii. Planar and Complete bipartite graph

**Ans.** Refer Q. 5.16.

Q. 6.

- a. Prove that a connected graph  $G$  is Euler graph if and only if every vertex of  $G$  is of even degree.
- b. Which of the following simple graph have a Hamiltonian circuit or, if not a Hamiltonian path ?



**Fig. 1.**

**Ans.** Refer Q. 5.17.

Q. 7. Prove that a simple graph with  $n$  vertices and  $k$  components can have at most  $\frac{(n-k)(n-k+1)}{2}$  edges.

**Ans.** Refer Q. 5.18.

Q. 8. What are different ways to represent a graph ? Define Euler circuit and Euler graph. Give necessary and sufficient conditions for Euler circuits and paths.

**Ans.** Refer Q. 5.19.

Q. 9. Define and explain any two the following :

1. BFS and DFS in trees
2. Euler graph
3. Adjacency matrix of a graph

**Ans.** Refer Q. 5.20.

Q. 10. Solve the recurrence relation  $y_{n+2} - 5y_{n+1} + 6y_n = 5^n$  subject to the condition  $y_0 = 0, y_1 = 2$ .

ANS. Refer Q. 5.24.

Q. 11. Solve the recurrence relation using generating function :

$$a_n - 7a_{n-1} + 10a_{n-2} = 0 \text{ with } a_0 = 3, a_1 = 3.$$

ANS. Refer Q. 5.25.

Q. 12. Solve the recurrence relation  $a_{r+2} - 5a_{r+1} + 6a_r = (r+1)^2$ .

ANS. Refer Q. 5.26.

Q. 13. Solve  $a_r - 6a_{r-1} + 8a_{r-2} = r \cdot 4^r$ , given  $a_0 = 8$ , and  $a_1 = 1$ .

ANS. Refer Q. 5.27.

Q. 14. Solve the recurrence relation :  $a_r + 4a_{r-1} + 4a_{r-2} = r^2$ .

ANS. Refer Q. 5.28.

Q. 15. Suppose that a valid codeword is an  $n$ -digit number in decimal notation containing an even number of 0's. Let  $a_n$  denote the number of valid codewords of length  $n$  satisfying the recurrence relation  $a_n = 8a_{n-1} + 10^{n-1}$  and the initial condition  $a_1 = 9$ . Use generating functions to find an explicit formula for  $a_n$ .

ANS. Refer Q. 5.29.

Q. 16. Suppose that a cookie shop has four different kinds of cookies. How many different ways can six cookies be chosen ?

ANS. Refer Q. 5.31.

Q. 17. A total of 1232 student have taken a course in Spanish, 879 have taken a course in French, and 114 have taken a course in Russian. Further 103 have taken courses in both Spanish and French, 23 have taken courses in both Spanish and Russian, and 14 have taken courses in both French and Russian. If 2092 students have taken least one of Spanish, French and Russian, how many students have taken a course in all three languages ?

ANS. Refer Q. 5.35.

